

Resultant as Determinant of Koszul Complex

A.Anokhina, A.Morozov and Sh.Shakirov¹

ITEP, Moscow, Russia
MIPT, Dolgoprudny, Russia

Abstract

A linear map between two vector spaces has a very important characteristic: a determinant. In modern theory two generalizations of linear maps are intensively used: to linear complexes (the nilpotent chains of linear maps) and to non-linear mappings. Accordingly, determinant of a linear map has two generalizations: to determinants of complexes and to resultants. These quantities are in fact related: resultant of a non-linear map is determinant of the corresponding Koszul complex. We give an elementary introduction into these notions and interrelations, which will definitely play a role in the future development of theoretical physics.

Contents

1	Introduction	2
2	Resultants	4
2.1	Properties of resultants	4
2.2	Examples of resultants	5
2.3	Sylvester method	6
2.4	Generalization of Sylvester's method: Koszul complex	7
3	Complexes and Determinants	10
3.1	Complex	10
3.2	Determinant of a complex	12
3.2.1	The case $1 \rightarrow 2 \rightarrow 1$	12
3.2.2	The case $2 \rightarrow 3 \rightarrow 1$	13
3.2.3	The case $1 \rightarrow 3 \rightarrow 2$	14
3.2.4	The case $2 \rightarrow 4 \rightarrow 2$	15
3.2.5	The case $n \rightarrow n + m \rightarrow m$	16
3.2.6	The case $n \rightarrow n + m \rightarrow m + k \rightarrow k$	17
3.2.7	The general case	19
3.3	Degree of determinant	20
3.4	How to calculate determinant of complex: concise summary	21
4	Koszul complexes	22
5	Evaluation of $R_{3 2}$	26
6	Conclusion	30

¹SashaAnokhina@yandex.ru; morozov@itep.ru; shakirov@itep.ru

1 Introduction

The origin of matrices and determinants lies deep in ancient civilizations. Around the second century BC, the Babylonians studied linear equations and the Chinese used matrices to find solutions to simple systems of linear equations, in two or three unknowns [1]. However, only in the late 17-th century these ideas became widely known and flourished with different applications. The theory of *linear* equations and transformations, now known as *linear algebra*, has completely proved its worth not only in pure science, but also in applied problems and in engineering. For example, Gauss himself applied linear algebra to determine the orbit of the asteroid Pallas [2]. Using observations of Pallas taken between 1803 and 1809, Gauss obtained and solved a system of six linear equations in six unknowns.

Soon after that, it became clear to researchers (Cayley, Sylvester and others [3, 4]) that linear algebra is just the simplest case of more general framework – of *polynomial* equations and transformations. They realized, that many parts of linear algebra have natural generalizations to arbitrary polynomials, not necessarily linear. Notably, determinant has such a generalization. A linear system in two unknowns

$$\begin{cases} f_0x + f_1y = 0 \\ g_0x + g_1y = 0 \end{cases}$$

is solvable (has a non-trivial solution x, y) if and only if its determinant vanishes

$$f_0g_1 - f_1g_0 = 0$$

and in complete analogy, a quadratic system

$$\begin{cases} f_0x^2 + f_1xy + f_2y^2 = 0 \\ g_0x^2 + g_1xy + g_2y^2 = 0 \end{cases}$$

is solvable if and only if certain expression, depending on coefficients, vanishes:

$$f_0^2g_2^2 - f_1f_2g_0g_1 + f_1^2g_0g_2 - 2f_0f_2g_0g_2 + f_0f_2g_1^2 - f_0f_1g_1g_2 + f_2^2g_0^2 = 0$$

This expression – a generalization of 2×2 determinant from linear to quadratic equations – is called a *resultant*. It was Sylvester who first studied it [4] and found the analogues for cubic, quartic and equations of higher degrees. Resultants also exist for more than two variables: in this case they provide non-linear generalizations of $n \times n$ determinant. A system

$$\begin{cases} f_1(x_1, x_2, \dots, x_n) = 0 \\ f_2(x_1, x_2, \dots, x_n) = 0 \\ \dots \\ f_n(x_1, x_2, \dots, x_n) = 0 \end{cases} \quad (1.1)$$

of n homogeneous polynomials of degree r in n variables

$$f_i(x_1, x_2, \dots, x_n) = \sum_{j_1, \dots, j_r=1}^n f_i^{j_1, j_2, \dots, j_r} x_{j_1} x_{j_2} \dots x_{j_r}$$

is solvable, if and only if an expression called its resultant vanishes:

$$R_{n|r}\{f_1, f_2, \dots, f_n\} = 0$$

For linear systems resultant $R_{n|1}$ is just a determinant, the well-known and well-understood object of linear algebra. Unfortunately, the knowledge and understanding of general resultants is not equally deep, yet. Moreover, the very direction of generalization – from linear algebra to arbitrary algebraic equations – has been almost completely forgotten after the 19-th century, and was revived only recently, after profound book of Gelfand, Kapranov and Zelevinsky [5].

For several reasons, we suggest to call this direction *non-linear algebra* [6]. It is expected to have as many applications as its linear cousin, if not more. Resultants – the main "special functions" of non-linear algebra – are already used in physics to describe, for example, multiparticle entanglement [7], singularities of Calabi-Yau three-folds [8], discrete dynamics [9]. There are applications to modeling and robotics [10]. However, applications remain restricted, partially because calculation of resultants is not as easy as that of determinants, especially when the number of variables is large.

It is often convenient to speak about systems of equations in terms of maps. Linear algebra studies linear maps between two spaces of dimensions m and n

$$L_1 \xrightarrow{f} L_2$$

which are given by linear polynomials:

$$f : \begin{pmatrix} x_1 \\ x_2 \\ \dots \\ x_n \end{pmatrix} \mapsto \begin{pmatrix} f_{11}x_1 + f_{12}x_2 + \dots + f_{1n}x_n \\ f_{21}x_1 + f_{22}x_2 + \dots + f_{2n}x_n \\ \dots \\ f_{m1}x_1 + f_{m2}x_2 + \dots + f_{mn}x_n \end{pmatrix}$$

Accordingly, non-linear algebra studies non-linear (polynomial) maps:

$$f : \begin{pmatrix} x_1 \\ x_2 \\ \dots \\ x_n \end{pmatrix} \mapsto \begin{pmatrix} f_1(x_1, x_2, \dots, x_n) \\ f_2(x_1, x_2, \dots, x_n) \\ \dots \\ f_m(x_1, x_2, \dots, x_n) \end{pmatrix} \quad (1.2)$$

System (1.1) corresponds to the particularly interesting case of $m = n$. When the system (1.1) is solvable, the corresponding map (1.2) is degenerate, i.e., has a non-vanishing kernel. As one can see, resultant defines a degeneracy condition for polynomial maps. Of course, it is very interesting to study further generalizations: to arbitrary non-linear mappings (not necessarily polynomial). For such generalizations and appropriate notion of resultant, see [11, 12].

The second, essentially different, generalization of determinant was introduced by Cayley approximately at the same time [3]. Instead of a single linear map between two spaces

$$L_1 \xrightarrow{f} L_2$$

he considered a *complex* – sequence of two linear maps between three spaces

$$L_1 \xrightarrow{f_1} L_2 \xrightarrow{f_2} L_3$$

subject to *nilpotency* constraint $f_2 \circ f_1 = 0$. Generally, the number of spaces can be arbitrary

$$L_1 \xrightarrow{f_1} L_2 \xrightarrow{f_2} L_3 \xrightarrow{f_3} \dots$$

and the maps satisfy $f_{i+1} \circ f_i = 0$ for all i . Cayley has found a quantity, associated to complexes in the same way as determinant is associated to linear maps. This quantity is now called *determinant of a complex*. For example, consider a sequence of two maps between spaces of dimensions 1, 2 and 1:

$$f_1 : x \mapsto \begin{pmatrix} a_1x \\ a_2x \end{pmatrix}, \quad f_2 : \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \mapsto b_1y_1 + b_2y_2$$

Composition of these maps

$$f_2 \circ f_1 : x \mapsto (a_1b_1 + a_2b_2)x$$

is requested to vanish. Cayley's determinant $\text{DET}(f_1, f_2)$ can be represented in three different ways

$$\text{DET}(f_1, f_2) = \frac{a_2}{b_1} = -\frac{a_1}{b_2} = \frac{|b|}{|a|}$$

where $|a|$ is simply the Euclidean length of vector $a = (a_1, a_2)$. Equivalence of all three follows from the above condition $a_1b_1 + a_2b_2 = 0$. This example shows, that determinant of complex is not a polynomial, it is a rational function. It vanishes when $|b| = 0$ and is singular when $|a| = 0$. It may be not obvious at the moment, but we will see below, that such a quantity is indeed a natural generalization of determinant

and has a number of important properties. In particular, determinant of a complex is (up to overall sign) invariant under permutations of basis vectors in the linear spaces.

It should be emphasized, that complexes are essentially objects of linear algebra and can be effectively treated using linear methods. For this reason the theory of complexes and their determinants, which is now called *homological algebra*, is more developed and attracts incomparably more attention, than resultant theory. Homological algebra has a wide range of physical applications, from topological field theories and invariants [13] to Faddeev-Popov's ghosts and Batalin-Vilkovisky approach to quantizing particle Lagrangians and Lagrangians of string field theory [14] and further to description of branes in terms of derived categories [15].

It has been known since Cayley, that resultants of non-linear maps and determinants of complexes are, in fact, related. Namely, given a non-linear map one can construct a complex, called *Koszul complex*, whose determinant is equal to resultant of the original map. Actually, this relation is one of a few reliable ways to calculate resultants, known at the moment. Several other ways to calculate resultants, not involving complexes, are described in [16, 11]. The aim of this paper is to explain this relation in clear and practical terms, making the text accessible for beginners. It can be considered as pure pedagogical, since most of the results are quite classical and known to the experts (see, e.g. [5, 17, 18]).

2 Resultants

2.1 Properties of resultants

Thus it is interesting and important to calculate resultants. Before discussing the calculation methods, we formulate the basic properties of resultants. As mentioned above, resultant $R_{n|r}\{f_1, f_2, \dots, f_n\}$ defines a degeneracy condition for polynomial maps of degree r in n variables

$$f : \begin{pmatrix} x_1 \\ x_2 \\ \dots \\ x_n \end{pmatrix} \mapsto \begin{pmatrix} f_1(x_1, x_2, \dots, x_n) \\ f_2(x_1, x_2, \dots, x_n) \\ \dots \\ f_n(x_1, x_2, \dots, x_n) \end{pmatrix}$$

where

$$f_i(x_1, x_2, \dots, x_n) = \sum_{j_1, \dots, j_r=1}^n f_i^{j_1, j_2, \dots, j_r} x_{j_1} x_{j_2} \dots x_{j_r}$$

Resultant $R_{n|r}(f)$ vanishes on degenerate maps and only on them. To say it another way, resultant is a common divisor of all polynomials, which vanish on degenerate maps. By this very definition, $R_{n|r}(f)$ is an irreducible polynomial in coefficients of f . Irreducibility of resultant is a generalization of the well-known fact, that determinant of a generic matrix is irreducible. For illustration, resultant

$$R_{3|1}(f) = \begin{vmatrix} f_{11} & f_{12} & f_{13} \\ f_{21} & f_{22} & f_{23} \\ f_{31} & f_{32} & f_{33} \end{vmatrix} = f_{11}f_{22}f_{33} - f_{11}f_{23}f_{32} - f_{21}f_{12}f_{33} + f_{21}f_{13}f_{32} + f_{31}f_{12}f_{23} - f_{31}f_{13}f_{22}$$

does not decompose into smaller factors. However, for non-generic maps – say, if some coefficients vanish – such decomposition does take place. For illustration, the resultant

$$\begin{vmatrix} f_{11} & 0 & 0 \\ f_{21} & f_{22} & f_{23} \\ f_{31} & f_{32} & f_{33} \end{vmatrix} = f_{11}(f_{22}f_{33} - f_{23}f_{32})$$

decomposes into smaller factors. The same holds for higher degrees of equations. For further study of resultant's irreducibility properties, see [19].

Another important property of $R_{n|r}$ is that it has a simple interpretation in terms of polynomials' roots. For example, consider a system of two polynomials of arbitrary degree in two variables:

$$f(x, y) = \prod_{i=1}^r (x - \alpha_i y)$$

$$g(x, y) = \prod_{i=1}^r (x - \beta_i y)$$

Resultant of this system is given by a simple formula

$$R_{2|r} \sim \prod_{i,j=1}^r (\beta_i - \alpha_j)$$

because the above system is solvable, if and only if $\alpha_j = \beta_i$ for some i, j . In fact, the right hand side is nothing but the product of values of $f(x, y)$ on 2-vectors $(\beta_1, 1) \dots (\beta_r, 1)$, which are roots of $g(x, y)$:

$$R_{2|r} \sim f(\vec{\beta}_1) \dots f(\vec{\beta}_r), \quad \vec{\beta}_i = \text{roots of } g(x, y) \quad (2.1)$$

Generalization to the case of n variables is straightforward. Resultant of a system

$$\begin{cases} f_1(x_1, \dots, x_n) = 0 \\ f_2(x_1, \dots, x_n) = 0 \\ \dots \\ f_n(x_1, \dots, x_n) = 0 \end{cases}$$

where $f_1, f_2, \dots, f_n$ are homogeneous polynomials of degree r , is given by a simple formula

$$R_{n|r} \sim f(\vec{\beta}_1) \dots f(\vec{\beta}_N), \quad \vec{\beta}_i = \text{common roots of } f_2(x_1, \dots, x_n), \dots, f_n(x_1, \dots, x_n) \quad (2.2)$$

The number N of common roots of a system of $(n - 1)$ equations of degrees $r_1, r_2, \dots, r_{n-1}$ is equal to

$$N = r_1 r_2 \dots r_{n-1}$$

In our case, we have all $r_i = r$, therefore the number of common roots is $N = r^{n-1}$. It immediately follows that $R_{n|r}$ is a homogeneous polynomial in coefficients of the first equation of degree N . Since (2.2) can be analogously written with any f_i in place of f_1 , resultant $R_{n|r}$ in fact has degree N in coefficients of any particular equation f_i . Consequently, it is a homogeneous polynomial of degree

$$\deg(R_{n|r}) = nN = nr^{n-1} \quad (2.3)$$

in coefficients of all equations. For $r = 1$, we recover the degree of determinant: $\deg R_{n|1} = n$. For systems of quadratic equations, $\deg R_{n|1} = 2^n n$, for cubic equations $\deg R_{n|1} = 3^n n$, and so on.

Despite being simple, the formula (2.2) – called *Poisson product formula* – is not very useful for calculation of resultants, because it is rather hard to find common roots of $(n - 1)$ non-linear equations in n variables. The roots are complicated functions of coefficients, which generally can not be expressed through radicals. Fortunately, Poisson product formula can be rewritten in several ways, which do not contain the roots explicitly. Such reformulations [16, 11] are more convenient in practical calculations.

2.2 Examples of resultants

Case $r = 1$. The simplest example is a linear map, for certainty in two variables:

$$f : \begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} f_0 x + f_1 y \\ g_0 x + g_1 y \end{pmatrix}$$

The map f is degenerate exactly when the linear system

$$\begin{cases} f_0 x + f_1 y = 0 \\ g_0 x + g_1 y = 0 \end{cases}$$

has a non-vanishing solution, i.e. when its determinant vanishes:

$$\begin{vmatrix} f_0 & f_1 \\ g_0 & g_1 \end{vmatrix} = 0$$

Therefore, resultant $R_{2|1}$ is nothing but a 2×2 determinant:

$$R_{2|1}(f) = \det_{2 \times 2}(f) = \begin{vmatrix} f_0 & f_1 \\ g_0 & g_1 \end{vmatrix}$$

In complete analogy, for linear maps in n variables resultant $R_{n|1}$ is just a $n \times n$ determinant:

$$R_{n|1}(f) = \det_{n \times n}(f) = \begin{vmatrix} f_{11} & f_{12} & \dots & f_{1n} \\ f_{21} & f_{22} & \dots & f_{2n} \\ \dots & \dots & \dots & \dots \\ f_{n1} & f_{n2} & \dots & f_{nn} \end{vmatrix}$$

Case $n = 2, r = 2$. As an example of non-linear maps, let us consider a quadratic map

$$f : \begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} f_0x^2 + f_1xy + f_2y^2 \\ g_0x^2 + g_1xy + g_2y^2 \end{pmatrix}$$

Degeneracy of f implies existence of non-zero x and y such that

$$\begin{cases} f_0x^2 + f_1xy + f_2y^2 = 0 \\ g_0x^2 + g_1xy + g_2y^2 = 0 \end{cases}$$

Provided that $f_0 \neq 0$ and $g_0 \neq 0$ the latter system can be written as:

$$\begin{cases} f_0(x - \alpha_1y)(x - \alpha_2y) = 0 \\ g_0(x - \beta_1y)(x - \beta_2y) = 0 \end{cases}$$

The system has non-trivial solution if and only if the polynomials have a common root, i.e. when

$$(\alpha_1 - \beta_1)(\alpha_1 - \beta_2)(\alpha_2 - \beta_1)(\alpha_2 - \beta_2) = 0$$

Since in this case explicit formulas for roots are known, one can express the resultant through the coefficients of f and g directly. After the expansion, we obtain a polynomial (if we include a multiplier $f_0^2g_0^2$):

$$\begin{aligned} R_{2|2}(f) &= f_0^2g_0^2(\alpha_1 - \beta_1)(\alpha_1 - \beta_2)(\alpha_2 - \beta_1)(\alpha_2 - \beta_2) = f_0^2(g_0^2\alpha_1^2 + g_1\alpha_1 + g_2)(g_0^2\alpha_2^2 + g_1\alpha_2 + g_2) = \\ &= f_0^2(g_0^2\alpha_1^2\alpha_2^2 + g_0g_1\alpha_1\alpha_2(\alpha_1 + \alpha_2) + g_0g_2((\alpha_1 + \alpha_2)^2 - 2\alpha_1\alpha_2) + g_1^2\alpha_1\alpha_2 + g_1g_2(\alpha_1 + \alpha_2) + g_2^2) = \\ &= g_0^2f_2^2 - g_0g_1f_1f_2 + g_0g_2(f_1^2 - 2f_0f_2) + g_1^2f_0f_2 - g_1g_2f_0f_1 + f_0^2g_2^2 \end{aligned}$$

In this way, we can calculate a resultant of two quadratic equations in two unknowns. Note, that it has degree 4 in map's coefficients, in accord with the general formula $nr^{n-1} = 2 \cdot 2^1 = 4$. However, it is hardly possible to use such an approach, when the degree of equations (or, worse than that, the number of variables) is greater than two. One needs another method to go beyond the simplest examples.

2.3 Sylvester method

The most classical resultant calculating method, which allows to go beyond the simplest examples, was found by Sylvester [4]. This method is to express resultant as a determinant of some auxiliary matrix. Such expression tautologically exists for linear maps in two variables

$$f : \begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} f_0x + f_1y \\ g_0x + g_1y \end{pmatrix} \quad R_{2|1}\{f, g\} = \begin{vmatrix} f_0 & f_1 \\ g_0 & g_1 \end{vmatrix} \quad (2.4)$$

and Sylvester has found, that it also exists for quadratic maps in two variables

$$f : \begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} f_0x^2 + f_1xy + f_2y^2 \\ g_0x^2 + g_1xy + g_2y^2 \end{pmatrix} \quad R_{2|2}\{f, g\} = \begin{vmatrix} f_0 & f_1 & f_2 & 0 \\ 0 & f_0 & f_1 & f_2 \\ g_0 & g_1 & g_2 & 0 \\ 0 & g_0 & g_1 & g_2 \end{vmatrix} \quad (2.5)$$

as well as for cubic maps in two variables

$$f : \begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} f_0x^3 + f_1x^2y + f_2xy^2 + f_3y^3 \\ g_0x^3 + g_1x^2y + g_2xy^2 + g_3y^3 \end{pmatrix}, \quad R_{2|3}\{f, g\} = \begin{vmatrix} f_0 & f_1 & f_2 & f_3 & 0 & 0 \\ 0 & f_0 & f_1 & f_2 & f_3 & 0 \\ 0 & 0 & f_0 & f_1 & f_2 & f_3 \\ g_0 & g_1 & g_2 & g_3 & 0 & 0 \\ 0 & g_0 & g_1 & g_2 & g_3 & 0 \\ 0 & 0 & g_0 & g_1 & g_2 & g_3 \end{vmatrix} \quad (2.6)$$

and for all higher degrees in two variables:

$$f : \begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} f_0x^r + f_1x^{r-1}y + \dots + f_ry^r \\ g_0x^r + g_1x^{r-1}y + \dots + g_ry^r \end{pmatrix}, \quad R_{2|r}(f) = \begin{vmatrix} f_1 & f_2 & \dots & f_r & 0 & 0 & \dots & 0 \\ 0 & f_1 & f_2 & \dots & f_r & 0 & \dots & 0 \\ \dots & 0 & \dots & 0 & f_1 & f_2 & \dots & f_r \\ g_0 & g_1 & \dots & g_r & 0 & 0 & \dots & 0 \\ 0 & g_0 & g_1 & \dots & g_r & 0 & \dots & 0 \\ \dots & 0 & \dots & 0 & g_0 & g_1 & \dots & g_r \end{vmatrix} \quad (2.7)$$

The beauty and simplicity made this formula and corresponding resultant $R_{2|r}$ widely known. Note, that it has degree $2r$ in map's coefficients, in accord with the general formula $nr^{n-1} = 2 \cdot r^1 = 2r$.

In our days, Sylvester method is the main apparatus used to calculate two-dimensional resultants and it is the only piece of resultant theory included in undergraduate text books (see for example [20]). However, its generalization to $n > 2$ is not straightforward. One possible generalization, representing resultant as *the determinant of Koszul complex*, is the subject of present paper. This generalization of Sylvester formula is the most widely known one. Other generalizations, which produce determinantal formulas for $n > 2$, are still the subject of active research (see, for example, [17]).

2.4 Generalization of Sylvester's method: Koszul complex

To describe Koszul complexes, we need several definitions. The basic object we deal with are homogeneous polynomials. All homogeneous polynomials $F^{i_1 \dots i_p} x_{i_1} \dots x_{i_p}$ of degree p in n variables form a linear space with a monomial basis $x_{i_1} \dots x_{i_p}$. Introduce now n auxiliary Grassmanian variables θ such, that

$$\theta_i \theta_j + \theta_j \theta_i = 0$$

The set of homogeneous polynomials of degree p in $x_1, \dots, x_n$ and of degree q in $\theta_1, \dots, \theta_n$ is again a linear space with monomial basis $x_{i_1} \dots x_{i_p} \theta_{j_1} \dots \theta_{j_q}$. We will use the notation $\Omega(p, q)$ for this linear space. Note that $q \leq n$: if a polynomial has degree in θ -variables bigger than n , it vanishes identically, because θ are anti-commuting variables. Introduction of such anti-commuting variables leads to a new possibility: given a polynomial map

$$x_j \rightarrow f_j(x) = f_j^{i_1 i_2 \dots i_r} x_{i_1} x_{i_2} \dots x_{i_r}$$

one can construct a linear operator

$$\hat{d} = f_j(x) \frac{\partial}{\partial \theta_j}$$

which is nilpotent:

$$\hat{d}\hat{d} = f_j(x) f_k(x) \frac{\partial}{\partial \theta_j} \frac{\partial}{\partial \theta_k} = f_j(x) f_k(x) \left(\frac{\partial}{\partial \theta_j} \frac{\partial}{\partial \theta_k} + \frac{\partial}{\partial \theta_k} \frac{\partial}{\partial \theta_j} \right) = 0$$

This operator, called *Koszul differential*, acts on the spaces $\Omega(p, q)$ in the following way:

$$\Omega(p, q) \xrightarrow{\hat{d}} \Omega(p+r, q-1) \xrightarrow{\hat{d}} \Omega(p+2r, q-2) \rightarrow \dots$$

In this way, operator $\hat{d}$ generates a sequence of linear spaces, called a *Koszul complex*. If we choose some basis in each linear space, operator $\hat{d}$ will be represented by a sequence of matrices (generally, not a single matrix). The idea – generalization of Sylvester's idea – is that resultant of f can be expressed through minors of these matrices. Before discussing the details, we illustrate this idea with simple examples.

The case of $n = 2, r = 1$. The simplest example is a linear map in two variables:

$$f : \begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} f_0x + f_1y \\ g_0x + g_1y \end{pmatrix}$$

By definition, differential is

$$\hat{d} = (f_0x + f_1y) \frac{\partial}{\partial \theta_1} + (g_0x + g_1y) \frac{\partial}{\partial \theta_2}$$

It acts on linear spaces

$$\Omega(1, 0) \xrightarrow{\hat{d}} \Omega(0, 1)$$

in the following way:

$$\begin{pmatrix} \theta_1 \\ \theta_2 \end{pmatrix} \xrightarrow{\hat{d}} \begin{pmatrix} f_0x + f_1y \\ g_0x + g_1y \end{pmatrix}$$

In the above pair of bases, differential is represented by a 2×2 matrix:

$$d = \begin{vmatrix} f_0 & f_1 \\ g_0 & g_1 \end{vmatrix}$$

and one can see that:

$$R_{2|1} = \det d = \begin{vmatrix} f_0 & f_1 \\ g_0 & g_1 \end{vmatrix}$$

This example illustrates the definition of Koszul differential. Of course, it is oversimplified, because the matrix of operator $\hat{d}$ in this particular case coincides with the matrix of map f .

The case of $n = 2, r = 2$. In this case the map is

$$f : \begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} f_0x^2 + f_1xy + f_2y^2 \\ g_0x^2 + g_1xy + g_2y^2 \end{pmatrix}$$

and the differential is

$$\hat{d} = (f_0x^2 + f_1xy + f_2y^2) \frac{\partial}{\partial \theta_1} + (g_0x^2 + g_1xy + g_2y^2) \frac{\partial}{\partial \theta_2}$$

It acts on linear spaces

$$\Omega(0, 1) \xrightarrow{\hat{d}} \Omega(2, 0)$$

in the following way:

$$\begin{pmatrix} \theta_1x \\ \theta_2x \\ \theta_1y \\ \theta_2y \end{pmatrix} \xrightarrow{\hat{d}} \begin{pmatrix} (f_0x^2 + f_1xy + f_2y^2)x \\ (g_0x^2 + g_1xy + g_2y^2)x \\ (f_0x^2 + f_1xy + f_2y^2)y \\ (g_0x^2 + g_1xy + g_2y^2)y \end{pmatrix}$$

In the above pair of bases, differential is represented by a 4×4 matrix:

$$d = \begin{vmatrix} f_0 & f_1 & f_2 & 0 \\ 0 & f_0 & f_1 & f_2 \\ g_0 & g_1 & g_2 & 0 \\ 0 & g_0 & g_1 & g_2 \end{vmatrix}$$

As one can see, we have obtained exactly the Sylvester's matrix. In accord with (2.5)

$$R_{2|2} = \det d = f_0^2g_2^2 - f_1f_2g_0g_1 + f_1^2g_0g_2 - 2f_0f_2g_0g_2 + f_0f_2g_1^2 - f_0f_1g_1g_2 + f_2^2g_0^2$$

And this is not an accident: if f has a non-trivial kernel, then the system

$$\begin{cases} (f_0x^2 + f_1xy + f_2y^2)x = 0 \\ (f_0x^2 + f_1xy + f_2y^2)y = 0 \\ (g_0x^2 + g_1xy + g_2y^2)x = 0 \\ (g_0x^2 + g_1xy + g_2y^2)y = 0 \end{cases}$$

has a non-vanishing solution and (since this system is linear in monomials x^3, x^2y, xy^2, y^3) its determinant $\det d$ must vanish. Therefore, if resultant vanishes, then determinant $\det d$ must vanish:

$$R_{2|2} = 0 \Rightarrow \det d = 0$$

For this reason

$$\det d = \mu R_{2|2}$$

Since both $\det d$ and $R_{2|2}$ are polynomials of degree 4 in f_0, f_1, g_0, g_1 , the coefficient μ is just a numeric factor. This example illustrates the use of Koszul differential in the case of non-linear maps.

The case of $n = 2, r = 1$: an example of 3-term complex. In both of the examples above, differential $\hat{d}$ was represented by a single square matrix. Now we will consider an example, where non-trivial complex – a sequence of several matrices – arises. We return to linear maps in two variables, where $\hat{d}$ has a form

$$\hat{d} = (f_0x + f_1y) \frac{\partial}{\partial \theta_1} + (g_0x + g_1y) \frac{\partial}{\partial \theta_2}$$

We will consider its action on another spaces

$$\Omega(0, 2) \xrightarrow{\hat{d}} \Omega(1, 1) \xrightarrow{\hat{d}} \Omega(2, 0)$$

In bases

$$(\theta_1\theta_2), \quad (\theta_1x, \theta_1y, \theta_2x, \theta_2y), \quad (x^2, xy, y^2)$$

differential is represented by a pair of rectangular matrices (not a single square matrix this time):

$$d_1 = \begin{vmatrix} -g_0 & -g_1 & f_0 & f_1 \end{vmatrix}, \quad d_2 = \begin{vmatrix} f_0 & f_1 & 0 \\ 0 & f_0 & f_1 \\ g_0 & g_1 & 0 \\ 0 & g_0 & g_1 \end{vmatrix}$$

Resultant $R_{2|1}$ is closely related to these matrices: the system with the matrix d_2

$$\begin{cases} (f_0x + f_1y)x = f_0x^2 + f_1xy + 0y^2 = 0 \\ (f_0x + f_1y)y = 0x^2 + f_0xy + f_1y^2 = 0 \\ (g_0x + g_1y)x = g_0x^2 + g_1xy + 0y^2 = 0 \\ (g_0x + g_1y)y = 0x^2 + g_0xy + g_1y^2 = 0 \end{cases}$$

which is a linear system in monomials x^2, xy, y^2 , has a non-zero solution if resultant $R_{2|1}$ vanishes. In this case, matrix d_2 becomes degenerate:

$$R_{2|1} = 0 \Rightarrow \text{all } 3 \times 3 \text{ minors of } d_2 \text{ are vanishing}$$

This is only possible, if all minors of d_2 are divisible by $R_{2|1}$. Therefore, $R_{2|1}$ should be a common factor of all these minors. As direct calculation shows

$$\begin{vmatrix} f_0 & f_1 & 0 \\ 0 & f_0 & f_1 \\ g_0 & g_1 & 0 \end{vmatrix} = -f_1 \begin{vmatrix} f_0 & f_1 \\ g_0 & g_1 \end{vmatrix}, \quad \begin{vmatrix} f_0 & f_1 & 0 \\ 0 & f_0 & f_1 \\ 0 & g_0 & g_1 \end{vmatrix} = f_0 \begin{vmatrix} f_0 & f_1 \\ g_0 & g_1 \end{vmatrix},$$

$$\begin{vmatrix} f_0 & f_1 & 0 \\ g_0 & g_1 & 0 \\ 0 & g_0 & g_1 \end{vmatrix} = g_1 \begin{vmatrix} f_0 & f_1 \\ g_0 & g_1 \end{vmatrix}, \quad \begin{vmatrix} 0 & f_0 & f_1 \\ g_0 & g_1 & 0 \\ 0 & g_0 & g_1 \end{vmatrix} = -g_0 \begin{vmatrix} f_0 & f_1 \\ g_0 & g_1 \end{vmatrix}$$

this is indeed true:

$$\text{common factor} = f_0g_1 - f_1g_0 = R_{2|1}$$

This example illustrates that, for one and the same polynomial map, there can be many different Koszul complexes, differing by the choice of linear spaces $\Omega(p, q)$. Depending on this choice, the complex can become either simpler or more complicated: in the above examples the simpler complex was

$$\Omega(1, 0) \xrightarrow{\hat{d}} \Omega(0, 1)$$

and the more complicated complex was

$$\Omega(0, 2) \xrightarrow{\hat{d}} \Omega(1, 1) \xrightarrow{\hat{d}} \Omega(2, 0)$$

In other words, the differential is fixed, but we are free to choose the linear spaces. In practice, one usually chooses a complex which is as simple as possible.

Moreover, this example illustrates the role of the first matrix d_1 . Since resultant is the common factor of minors of d_2 , we can determine resultant from the matrix d_2 only. From this point of view, it seems that d_1 does not provide any information about resultant. This is true, but only in part: d_1 provides precise information about the extraneous factors. As one can see, the other factors of minors d_2 are, up to a sign, elements of d_1 . Therefore, $R_{2|1}$ can be found by division of minors of d_2 on 'conjugate' minors of d_1 . This ratio is a particular example of what we call determinant of a complex. Remarkably, for Koszul complex the ratio is actually a *polynomial* in the coefficients of the map f .

3 Complexes and Determinants

3.1 Complex

A sequence of linear maps between several linear spaces

$$0 \xrightarrow{d_0} L_1 \xrightarrow{d_1} \dots \xrightarrow{d_{p-1}} L_p \xrightarrow{d_p} 0$$

is called a *p-term complex* iff nilpotency condition

$$\forall i \quad d_{i+1} \circ d_i = 0$$

is satisfied. Linear spaces L_i of dimensions $l_i = \dim L_i$ are called *terms* of the complex. Therefore, an ordinary linear map is a 2-term complex (a complex, which consists of two linear spaces and a single map between them). Operators $\hat{d}_i$ are called differential operators. According to the standard terminology, if a vector is equal to differential of some other vector (i.e, lies in the image of differential $\hat{d}_i$)

$$v = \hat{d}_i w$$

it is called *exact*. If a vector is annihilated by differential (i.e, lies in the kernel of $\hat{d}_{i+1}$)

$$\hat{d}_{i+1} v = 0$$

it is called *closed*. Equivalently, nilpotency condition can be expressed as

$$\forall i \quad \text{Im } d_i \subset \text{Ker } d_{i+1}$$

that is, any exact vector is closed. It is very important, that the inverse is, generally speaking, wrong: there can be vectors, which are closed but not exact. In other words, there can be vectors from $\text{Ker } d_{i+1}$ which do not belong to $\text{Im } d_i$. Such vectors again form a linear space

$$H_i = \text{Ker } d_{i+1} / \text{Im } d_i$$

of dimension $h_i = \dim H_i$, called *i-th cohomology space* of d . Elements of H_i are called cohomologies. A complex, which has no cohomologies (i.e, with $h_i = 0$ for all i) is called *exact complex*, because all closed elements in such a complex are exact. The notion of cohomology is so important in theory of complexes, that the whole field is known as "homological" algebra.




Figure 1: A schematic picture of a sequence of linear maps between several linear spaces.

It is evident, that $\dim (\text{Im } d_i) + \dim (\text{Ker } d_i) = \dim L_i$

It is well known from linear algebra, that

$$\dim (\text{Im } \hat{d}_i) + (\dim \text{Ker } \hat{d}_i) = \dim L_i$$

At the same time, by definition of cohomology

$$\dim (\text{Im } \hat{d}_i) = \dim (\text{Ker } \hat{d}_{i+1}) - \dim H_i$$

From these two identities, it follows

$$\dim L_i = \dim (\text{Ker } \hat{d}_i) + \dim (\text{Ker } \hat{d}_{i+1}) + \dim H_i$$

If we take an alternating sum, dimensions of the kernel spaces drop away:

$$\sum_{k=1}^p (-1)^k \dim L_k = \sum_{k=1}^p (-1)^k \dim H_k \equiv \chi \quad (3.1)$$

This alternating combination of dimensions is usually called *Euler characteristic* of the complex and denoted χ . As follows from (3.1), if the complex is exact, its Euler characteristic is zero:

$$\text{complex is exact} \Rightarrow l_1 - l_2 + l_3 - \dots + (-1)^p l_p = 0 \quad (3.2)$$

Inverse is, generally, wrong: vanishing Euler characteristic does not imply all $h_i = 0$, it implies only that

$$h_1 - h_2 + h_3 - \dots + (-1)^p h_p = 0$$

Consequently, (3.2) is a necessary condition for a complex to be exact. To say it in other words, if $\chi \neq 0$, the complex is definitely not exact, but if $\chi = 0$, we need some additional information to decide, whether a given complex is exact or not. This additional information is provided by determinant of the complex: determinant is non-vanishing only for exact complexes.

3.2 Determinant of a complex

In this section we derive an expression for determinant of a complex. We start from concrete examples, using vector notations and geometric analogies to simplify understanding. The aim is to create an intuitive image of what determinant of a complex is: a coefficient of proportionality between the two natural geometric quantities. Having this in mind, we will move from simple examples to general cases.

3.2.1 The case $1 \rightarrow 2 \rightarrow 1$

This simplest example is a 3-term complex $L_1 \xrightarrow{d_1} L_2 \xrightarrow{d_2} L_3$ with dimensions of linear spaces $l = (1, 2, 1)$. Its Euler characteristic is equal to $1 - 2 + 1 = 0$. The first differential sends

$$\hat{d}_1 : x \mapsto \vec{a}x$$

where x is a scalar from L_1 and $\vec{a}$ is some constant 2-vector from L_2 . The second differential sends

$$\hat{d}_2 : \vec{y} \mapsto \vec{b}\vec{y}$$

where $\vec{y}$ and $\vec{b}$ are 2-vectors from L_2 , and $\vec{b}\vec{y} = b_1y_1 + b_2y_2$ is a scalar product. As one can see, in this example linear maps d_1 and d_2 are parametrized by vectors $\vec{a}$ and $\vec{b}$, respectively. Their composition is

$$\hat{d}_2 \circ \hat{d}_1 : x \mapsto \vec{a}\vec{b} \cdot x$$

and the nilpotency condition is that $\vec{a}\vec{b} = 0$, i.e., $\vec{b}$ is orthogonal to $\vec{a}$. In two dimensions, all vectors orthogonal to $\vec{a}$ are proportional to $\epsilon_{ij}a_j$, i.e., to $(a_2, -a_1)$. Vector $\vec{b}$ should be also proportional to it:

$$b_i = \eta \cdot \epsilon_{ij}a_j$$

The coefficient of proportionality η is exactly the determinant of this complex. When $\eta \neq 0$, the kernel $\text{Ker } \hat{d}_2$ is 1-dimensional and consists of vectors, proportional to $\vec{a}$. It coincides with the image $\text{Im } \hat{d}_1$, so there are no cohomologies and complex is exact. When η vanishes, the kernel $\text{Ker } \hat{d}_2$ becomes 2-dimensional and the image $\text{Im } \hat{d}_1$ stays 1-dimensional, so $2 - 1 = 1$ -dimensional cohomologies appear and complex fails to be exact. Thus, η indeed allows to distinguish between exact and non-exact complexes.




Figure 2: Every vector in 2d, orthogonal to (a_1, a_2) , is proportional to $(a_2, -a_1)$

Determinant can be written through the components in 2 different ways:

$$\eta = \frac{b_1}{a_2}$$

$$\eta = -\frac{b_2}{a_1}$$

In the third, most symmetric form, determinant is equal to the ratio of lengths

$$\eta = \frac{|b|}{|a|} = \sqrt{\frac{b_1^2 + b_2^2}{a_1^2 + a_2^2}}$$

which emphasizes its orthogonal invariance. Equivalence of these ways follows from $a_1b_1 + a_2b_2 = 0$.

3.2.2 The case $2 \rightarrow 3 \rightarrow 1$

The next-to-simplest example is a 3-term complex $L_1 \xrightarrow{d_1} L_2 \xrightarrow{d_2} L_3$ with dimensions of linear spaces $l = (2, 3, 1)$. Its Euler characteristic is equal to $2 - 3 + 1 = 0$. The first differential sends

$$\hat{d}_1 : \vec{x} \mapsto \vec{a}x_1 + \vec{b}x_2$$

where $\vec{x}$ is a 2-vector from L_1 and $\vec{a}, \vec{b}$ is a pair of 3-vectors from L_2 . The second differential sends

$$\hat{d}_2 : \vec{y} \mapsto \vec{c}\vec{y}$$

where $\vec{y}$ and $\vec{c}$ are 3-vectors from L_2 , and $\vec{c}\vec{y} = c_1y_1 + c_2y_2 + c_3y_3$ is a scalar product. As one can see, in this example linear maps d_1 and d_2 are parametrized by $\vec{a}, \vec{b}$ and $\vec{c}$, respectively. Their composition is

$$\hat{d}_2 \circ \hat{d}_1 : \vec{x} \mapsto \vec{a}\vec{c} \cdot x_1 + \vec{b}\vec{c} \cdot x_2$$

with nilpotency condition

$$\vec{a}\vec{c} = \vec{b}\vec{c} = 0$$

i.e, $\vec{c}$ is orthogonal both to $\vec{a}$ and to $\vec{b}$. In three dimensions, all vectors orthogonal both to $\vec{a}$ and to $\vec{b}$ are proportional to $\epsilon_{ijk}a_jb_k$, i.e, to vector product $\vec{a} \times \vec{b}$. Vector $\vec{c}$ should be also proportional to it:

$$\vec{c} = \eta \cdot \vec{a} \times \vec{b}$$

$$c_i = \eta \cdot \epsilon_{ijk}a_jb_k$$

The coefficient of proportionality η is exactly the determinant of this complex. When $\eta \neq 0$, the kernel $\text{Ker } \hat{d}_2$ is 2-dimensional and consists of linear combinations of $\vec{a}$ and $\vec{b}$. It coincides with the image $\text{Im } \hat{d}_1$, so there are no cohomologies and complex is exact. When η vanishes, the kernel $\text{Ker } \hat{d}_2$ becomes 3-dimensional and the image $\text{Im } \hat{d}_1$ stays 2-dimensional, so $3 - 2 = 1$ -dimensional cohomologies appear and complex fails to be exact. Thus, η indeed allows to distinguish between exact and non-exact complexes.




Figure 3: Every vector in 3d, orthogonal both to $\vec{a}$ and to $\vec{b}$, is proportional to $\vec{a} \times \vec{b}$

Determinant can be written through the components in 3 different ways:

$$\eta = \frac{c_1}{a_2b_3 - a_3b_2}$$

$$\eta = \frac{c_2}{a_3b_1 - a_1b_3}$$

$$\eta = \frac{c_3}{a_1b_2 - a_2b_1}$$

Equivalence of these three representations follows from the nilpotency condition

$$a_1c_1 + a_2c_2 + a_3c_3 = b_1c_1 + b_2c_2 + b_3c_3 = 0$$

3.2.3 The case $1 \rightarrow 3 \rightarrow 2$

Our next example is a 3-term complex $L_1 \xrightarrow{d_1} L_2 \xrightarrow{d_2} L_3$ with dimensions of linear spaces $l = (1, 3, 2)$. Its Euler characteristic is equal to $1 - 3 + 2 = 0$. The first differential sends

$$\hat{d}_1 : x \mapsto \vec{a}x$$

where x is a scalar from L_1 and $\vec{a}$ is some constant 3-vector from L_2 . The second differential sends

$$\hat{d}_2 : \vec{y} \mapsto \begin{pmatrix} \vec{b}\vec{y} \\ \vec{c}\vec{y} \end{pmatrix}$$

where $\vec{y}, \vec{b}$ and $\vec{c}$ are 3-vectors from L_2 and the right hand side is a 2-vector from L_3 . As one can see, in this example linear maps d_1 and d_2 are parameterized by $\vec{a}$ and $\vec{b}, \vec{c}$, respectively. Their composition is

$$\hat{d}_2 \circ \hat{d}_1 : x \mapsto \begin{pmatrix} \vec{a}\vec{b} \\ \vec{a}\vec{c} \end{pmatrix} \cdot x$$

with nilpotency condition

$$\vec{a}\vec{b} = \vec{a}\vec{c} = 0$$

i.e, $\vec{a}$ is orthogonal both to $\vec{b}$ and to $\vec{c}$. This implies, in complete analogue with the previous example, that the vector product $\vec{b} \times \vec{c}$ is proportional to $\vec{a}$:

$$\vec{b} \times \vec{c} = \eta \cdot \vec{a}$$

$$\epsilon_{ijk} b_j c_k = \eta \cdot a_i$$

The coefficient of proportionality η is exactly the determinant of this complex. When $\eta \neq 0$, the kernel $\text{Ker } \hat{d}_2$ is 1-dimensional and consists of vectors, proportional to $\vec{a}$. It coincides with the image $\text{Im } \hat{d}_1$, so there are no cohomologies and complex is exact. When η vanishes, $\vec{b}$ and $\vec{c}$ become collinear and the kernel $\text{Ker } \hat{d}_2$ – by definition, the space of all vectors, orthogonal to $\vec{b}$ and $\vec{c}$ – becomes a 2-dimensional plane. The image $\text{Im } \hat{d}_1$ stays 1-dimensional, so $2 - 1 = 1$ -dimensional cohomologies appear and complex fails to be exact. Thus, η indeed allows to distinguish between exact and non-exact complexes.




Figure 4: The space of vectors, orthogonal to $\vec{b}$ and $\vec{c}$, is 1-dimensional when $\vec{b}$ and $\vec{c}$ are distinct.
The space of vectors, orthogonal to $\vec{b}$ and $\vec{c}$, is 2-dimensional when $\vec{b}$ and $\vec{c}$ are collinear.

Determinant can be written through the components in 3 different ways:

$$\eta = \frac{b_2 c_3 - b_3 c_2}{a_1}$$

$$\eta = \frac{b_3 c_1 - b_1 c_3}{a_2}$$

$$\eta = \frac{b_1 c_2 - b_2 c_1}{a_3}$$

Again, equivalence of these three representations follows from nilpotency conditions

$$a_1 b_1 + a_2 b_2 + a_3 b_3 = a_1 c_1 + a_2 c_2 + a_3 c_3 = 0$$

3.2.4 The case $2 \rightarrow 4 \rightarrow 2$

More complicated example is a 3-term complex $L_1 \xrightarrow{d_1} L_2 \xrightarrow{d_2} L_3$ with dimensions of linear spaces $l = (2, 4, 2)$. Its Euler characteristic is equal to $2 - 4 + 2 = 0$. The first differential sends

$$\hat{d}_1 : \vec{x} \mapsto \vec{a}x_1 + \vec{b}x_2$$

where x is a 2-vector from L_1 and $\vec{a}, \vec{b}$ is a pair of 4-vectors from L_2 . The second differential sends

$$\hat{d}_2 : \vec{y} \mapsto \begin{pmatrix} \vec{p}\vec{y} \\ \vec{q}\vec{y} \end{pmatrix}$$

where $\vec{y}, \vec{p}$ and $\vec{q}$ are 4-vectors from L_2 and the right hand side is a 2-vector from L_3 . As one can see, in this example linear maps d_1 and d_2 are parametrized by $\vec{a}, \vec{b}$ and $\vec{p}, \vec{q}$, respectively. Their composition is

$$\hat{d}_2 \circ \hat{d}_1 : \vec{x} \mapsto \begin{pmatrix} \vec{p}\vec{a} \\ \vec{q}\vec{a} \end{pmatrix} \cdot x_1 + \begin{pmatrix} \vec{p}\vec{b} \\ \vec{q}\vec{b} \end{pmatrix} \cdot x_2$$

with nilpotency condition

$$\vec{a}\vec{p} = \vec{a}\vec{q} = \vec{b}\vec{p} = \vec{b}\vec{q} = 0$$

Such constraints are quite typical for calculations with 4-vectors. Notice, that

$$p_\mu q_\nu - q_\nu p_\mu$$

is antisymmetric in μ, ν and satisfies relations

$$(p_\mu q_\nu - p_\nu q_\mu) a_\mu = 0$$

$$(p_\mu q_\nu - p_\nu q_\mu) b_\mu = 0$$

due to nilpotency. Another antisymmetric combination, which satisfies the same relations, is

$$\epsilon_{\mu\nu\alpha\beta} a_\alpha b_\beta$$

due to complete antisymmetry of the ϵ -tensor. Consequently, one must be proportional to another:

$$p_\mu q_\nu - q_\nu p_\mu = \eta \cdot \epsilon_{\mu\nu\alpha\beta} a_\alpha b_\beta$$

The coefficient of proportionality η is exactly the determinant of this complex. When $\eta \neq 0$, the kernel $\text{Ker } \hat{d}_2$ is 2-dimensional and consists of linear combinations of $\vec{a}$ and $\vec{b}$. It coincides with the image $\text{Im } \hat{d}_1$, so there are no cohomologies and complex is exact. When η vanishes, $\vec{p}$ and $\vec{q}$ become collinear and the kernel $\text{Ker } \hat{d}_2$ – by definition, the space of all vectors, orthogonal to $\vec{p}$ and $\vec{q}$ – becomes a 3-dimensional (hyper)plane. The image $\text{Im } \hat{d}_1$ stays 2-dimensional, so $3 - 2 = 1$ -dimensional cohomologies appear and complex fails to be exact. Thus, η indeed allows to distinguish between exact and non-exact complexes.

Determinant can be written through the components in 6 different ways:

$$\eta = \frac{p_1 q_2 - p_2 q_1}{a_3 b_4 - a_4 b_3}, \quad \eta = \frac{p_1 q_3 - p_3 q_1}{a_2 b_4 - a_4 b_2}$$

$$\eta = \frac{p_1 q_4 - p_4 q_1}{a_2 b_3 - a_3 b_2}, \quad \eta = \frac{p_2 q_3 - p_3 q_2}{a_1 b_4 - a_4 b_1}$$

$$\eta = \frac{p_2 q_4 - p_4 q_2}{a_1 b_3 - a_3 b_1}, \quad \eta = \frac{p_3 q_4 - p_4 q_3}{a_1 b_2 - a_2 b_1}$$

As usual equivalence of all these representations follows from the nilpotency conditions. We do not include a visualization here: a picture of orthogonal vectors in four dimensions is not very illuminating.




Figure 5: Choosing of minors in the case $2 \rightarrow 5 \rightarrow 3$.

Determinant is equal to the ratio of two "white" minors: $\frac{\text{Minor}_{3 \times 3}}{\text{Minor}_{2 \times 2}}$.

3.2.5 The case $n \rightarrow n + m \rightarrow m$

Summarizing the above examples, consider a 3-term complex $L_1 \xrightarrow{d_1} L_2 \xrightarrow{d_2} L_3$ with dimensions of linear spaces $l = (n, n + m, m)$. Its Euler characteristic is $n - n - m + m = 0$. The first differential sends

$$\hat{d}_1 : x_i \mapsto A_{ij} x_j$$

where A_{ij} is a $n \times (n + m)$ matrix, i.e, index i runs from 1 to n and index j runs from 1 to $(n + m)$. The second differential sends

$$\hat{d}_2 : y_i \mapsto B_{ij} y_j$$

where B_{ij} is a $(n + m) \times m$ matrix, i.e, index i runs from 1 to $(n + m)$ and index j runs from 1 to m . Vector notations, used in the previous examples, are convenient only in low dimensions and we will not use them anymore. From now on, we switch to matrix notations: as one can see, linear maps d_1 and d_2 are parametrized by matrices A and B . Their composition is

$$\hat{d}_2 \circ \hat{d}_1 : x_i \mapsto A_{ij} B_{jk} x_k$$

with nilpotency condition

$$A_{ij} B_{jk} = 0$$

Introduce the minors of matrices A, B :

$$(M_A)_{i_1 \dots i_n} = n \times n \text{ minor of } A, \text{ situated in the columns } i_1, \dots, i_n$$

$$(M_B)_{i_1 \dots i_n} = m \times m \text{ minor of } B, \text{ obtained by throwing away the rows } i_1, \dots, i_n$$

They are given by explicit expressions with the ϵ -tensors:

$$(M_A)_{i_1 \dots i_n} = \epsilon_{j_1 \dots j_n} A_{j_1 i_1} \dots A_{j_n i_n}$$

$$(M_B)_{i_1 \dots i_n} = \epsilon_{i_1 \dots i_n j_1 \dots j_m} \epsilon_{k_1 \dots k_m} B_{j_1 k_1} \dots B_{j_m k_m}$$

Notice, that M_A, M_B are both antisymmetric in $i_1, \dots, i_n$ and vanish when contracted with B :

$$(M_A)_{i_1 \dots i_n} \cdot B_{i_1 s_1} = 0$$

$$(M_B)_{i_1 \dots i_n} \cdot B_{i_1 s_1} = 0$$




Figure 6: Choosing of minors in the case $3 \rightarrow 9 \rightarrow 7 \rightarrow 1$. Determinant is equal to $\frac{\text{Minor}_{6 \times 6}}{\text{Minor}_{3 \times 3} \text{Minor}_{1 \times 1}}$

Tensor M_A does so due to nilpotency conditions, and M_B due to complete antisymmetry of the ϵ -tensor. Consequently, just like it was in the previous example, one must be proportional to another

$$(M_B)_{i_1 \dots i_n} = \eta \cdot (M_A)_{i_1 \dots i_n}$$

since it is the only way to satisfy such a restrictive set of conditions. In components we have

$$\epsilon_{i_1 \dots i_n j_1 \dots j_m} \epsilon_{k_1 \dots k_m} B_{j_1 k_1} \dots B_{j_m k_m} = \eta \cdot \epsilon_{j_1 \dots j_n} A_{j_1 i_1} \dots A_{j_n i_n}$$

The coefficient of proportionality η is exactly the determinant of this complex. When $\eta \neq 0$, the kernel $\text{Ker } \hat{d}_2$ is n -dimensional. It coincides with the image $\text{Im } \hat{d}_1$, so there are no cohomologies and complex is exact. When η vanishes, all minors of B vanish, so the kernel $\text{Ker } \hat{d}_2$ becomes $n + 1$ dimensional. The image $\text{Im } \hat{d}_1$ stays n -dimensional, so 1-dimensional cohomologies appear and complex fails to be exact. Thus, η indeed allows to distinguish between exact and non-exact complexes.

Determinant can be written through components in $C_{n+m}^n = (n+m)!/n!/m!$ different ways:

$$\eta = \text{DET}(A, B) = \frac{(M_B)_{i_1 \dots i_n}}{(M_A)_{i_1 \dots i_n}}$$

that is the number of components of antisymmetric tensor with n indices running from 1 to $n + m$. In other words, that is the number of ways to choose n items $i_1, \dots, i_n$ out of $n + m$ possibilities. Equivalence of these ways, as we have shown, follows from the nilpotency conditions. On the practical level, different ways correspond to different choices of rows and columns (an illustration is shown on Fig. 5).

3.2.6 The case $n \rightarrow n + m \rightarrow m + k \rightarrow k$

Moving further, consider a 4-term complex $L_1 \xrightarrow{d_1} L_2 \xrightarrow{d_2} L_3 \xrightarrow{d_3} L_4$ with dimensions of linear spaces $l = (n, n + m, m + k, k)$. Its Euler characteristic is $n - n - m + m + k - k = 0$. The first differential sends

$$\hat{d}_1 : x_i \mapsto A_{ij} x_j$$

where A_{ij} is a $n \times (n + m)$ matrix, i.e, index i runs from 1 to n and index j runs from 1 to $(n + m)$. The second differential sends

$$\hat{d}_2 : y_i \mapsto B_{ij} y_j$$

where B_{ij} is a $(n + m) \times (m + k)$ matrix, i.e, index i runs from 1 to $(n + m)$ and index j runs from 1 to $(m + k)$. The third differential sends

$$\hat{d}_3 : z_i \mapsto C_{ij} z_j$$

where C_{ij} is a $(m+k) \times k$ matrix, i.e, index i runs from 1 to $(m+k)$ and index j runs from 1 to k . Linear maps d_1, d_2, d_3 are parametrized by matrices A, B and C . Their compositions are equal to

$$\hat{d}_2 \circ \hat{d}_1 : x_i \mapsto A_{ij} B_{jk} x_k$$

$$\hat{d}_3 \circ \hat{d}_2 : y_i \mapsto B_{ij} C_{jk} y_k$$

with nilpotency conditions

$$A_{ij} B_{jk} = 0$$

$$B_{ij} C_{jk} = 0$$

The relevant minors of matrices A, B, C are

$$\begin{aligned} (M_A)_{i_1 \dots i_n} &= n \times n \text{ minor of } A, \text{ situated in columns } i_1, \dots, i_n \\ (M_B)_{i_1 \dots i_n | j_1 \dots j_m} &= m \times m \text{ minor of } B, \text{ obtained by crossing out rows } i_1, \dots, i_n \\ &\quad \text{and selecting columns } j_1, \dots, j_m \\ (M_C)_{j_1 \dots j_m} &= k \times k \text{ minor of } C, \text{ obtained by crossing out rows } j_1, \dots, j_m \end{aligned}$$

They are given by explicit expressions with ϵ -tensors:

$$\begin{aligned} (M_A)_{i_1 \dots i_n} &= \epsilon_{j_1 \dots j_n} A_{j_1 i_1} \dots A_{j_n i_n} \\ (M_B)_{i_1 \dots i_n | j_1 \dots j_m} &= \epsilon_{i_1 \dots i_n s_1 \dots s_m} B_{s_1 j_1} \dots B_{s_m j_m} \\ (M_C)_{j_1 \dots j_m} &= \epsilon_{j_1 \dots j_m s_1 \dots s_k} \epsilon_{w_1 \dots w_k} C_{s_1 w_1} \dots C_{s_k w_k} \end{aligned}$$

Let us show, that

$$(M_A)_{i_1 \dots i_n} (M_C)_{j_1 \dots j_m} = \eta \cdot (M_B)_{i_1 \dots i_n | j_1 \dots j_m}$$

This is easy: both expressions vanish when contracted with B

$$(M_A)_{i_1 \dots i_n} (M_C)_{j_1 \dots j_m} \cdot B_{i_1 s_1} = (M_B)_{i_1 \dots i_n | j_1 \dots j_m} \cdot B_{i_1 s_1} = 0$$

vanish when contracted with C

$$(M_A)_{i_1 \dots i_n} (M_C)_{j_1 \dots j_m} \cdot C_{j_1 s_1} = (M_B)_{i_1 \dots i_n | j_1 \dots j_m} \cdot C_{j_1 s_1} = 0$$

and both are antisymmetric separately in $i_1 \dots i_n$ and in $j_1 \dots j_m$. Consequently, they are proportional, since it is the only way to satisfy such a restrictive set of conditions.

The coefficient of proportionality η is exactly the determinant of this complex. When $\eta \neq 0$, matrices A and C have the full rank, what assures that there is no cohomology. When η vanishes, either all minors of A vanish or all minors of C vanish, cohomologies appear either in the first or in the third term, and complex fails to be exact. Thus, η indeed allows to distinguish between exact and non-exact complexes.

Determinant can be written through components in $C_{n+m}^n C_{m+k}^m$ different ways:

$$\eta = \text{DET } (A, B, C) = \frac{(M_A)_{i_1 \dots i_n} (M_C)_{j_1 \dots j_m}}{(M_B)_{i_1 \dots i_n | j_1 \dots j_m}}$$

that is the number of ways to choose n items $i_1, \dots, i_n$ out of $n + m$ possibilities *and* m items $j_1, \dots, j_m$ out of $m + k$ possibilities. Equivalence of these ways, as we have shown, follows from the nilpotency conditions. On the practical level, different ways correspond to different choices of rows and columns (an illustration is shown on Fig. 6).

3.2.7 The general case

Finally, consider a p -term complex $L_1 \xrightarrow{d_1} L_2 \xrightarrow{d_2} \dots L_{p-1} \xrightarrow{d_{p-1}} L_p$ with dimensions of linear spaces

$$l_m = k_{m-1} + k_m$$

where k_0 and k_p are by definition zero. Its Euler characteristic is equal to

$$\chi = \sum_{m=1}^p (-1)^{m-1} (k_{m-1} + k_m) = k_0 + (-1)^{p-1} k_p = 0$$

In fact, it is the general form of a complex with vanishing Euler characteristic. Differentials send

$$\hat{d}_m : x_i \mapsto (d_m)_{ij} x_j$$

where $(d_m)_{ij}$ is a $l_m \times l_{m+1}$ matrix, i.e. the index i runs from 1 to l_m and index j runs from 1 to l_{m+1} . In this way our complex is represented by a sequence of rectangular matrices. Compositions are equal to

$$\hat{d}_{m+1} \circ \hat{d}_m : x_i \mapsto (d_m)_{ij} (d_{m+1})_{jk} x_k$$

with nilpotency conditions

$$(d_m)_{ij} (d_{m+1})_{jk} = 0$$

Just like we did in our previous examples, consider the minors of matrices $d_1, \dots, d_{p-1}$:

$$\begin{aligned} (M_1)_{\sigma_2} &= k_1 \times k_1 \text{ minor of } d_1, \text{ situated in columns } \sigma_2 \\ (M_2)_{\sigma_2 | \sigma_3} &= k_2 \times k_2 \text{ minor of } d_2, \text{ obtained by crossing out rows } \sigma_2 \text{ and selecting columns } \sigma_3 \\ (M_3)_{\sigma_3 | \sigma_4} &= k_3 \times k_3 \text{ minor of } d_3, \text{ obtained by crossing out rows } \sigma_3 \text{ and selecting columns } \sigma_4 \\ &\dots\dots\dots \\ (M_i)_{\sigma_i | \sigma_{i+1}} &= k_i \times k_i \text{ minor of } d_i, \text{ obtained by crossing out rows } \sigma_i \text{ and selecting columns } \sigma_{i+1} \\ &\dots\dots\dots \\ (M_{p-1})_{\sigma_{p-1}} &= k_{p-1} \times k_{p-1} \text{ minor of } d_{p-1}, \text{ obtained by crossing out rows } \sigma_{p-1} \end{aligned}$$

where each $\sigma_i \subset \{1, 2, \dots, l_i\}$ is an arbitrary set of indices of length $|\sigma_i| = k_{i-1}$. They are given by

explicit (although a little unwieldy) expressions with ϵ -tensors, just like in the previous examples:

$$\begin{aligned} (M_1)_{a_1 \dots a_{k_1}} &= \epsilon_{s_1 \dots s_{k_1}} (d_1)_{s_1 a_1} \dots (d_1)_{s_{k_1} a_{k_1}} \\ (M_i)_{a_1 \dots a_{k_{i-1}} | b_1 \dots b_{k_i}} &= \epsilon_{a_1 \dots a_{k_{i-1}} s_1 \dots s_{k_i}} (d_i)_{s_1 b_1} \dots (d_i)_{s_{k_i} b_{k_i}} \\ (M_{p-1})_{a_1 \dots a_{k_{p-2}}} &= \epsilon_{a_1 \dots a_{k_{p-2}} s_1 \dots s_{k_{p-1}}} \epsilon_{j_1 \dots j_{k_{p-1}}} (d_{p-1})_{s_1 b_1} \dots (d_{p-1})_{s_{k_{p-1}} j_{k_{p-1}}} \end{aligned}$$

Let us show, that

$$(M_1)_{\sigma_2} (M_3)_{\sigma_3 | \sigma_4} (M_5)_{\sigma_5 | \sigma_6} \dots = \eta^{\pm 1} \cdot (M_2)_{\sigma_2 | \sigma_3} (M_4)_{\sigma_4 | \sigma_5} (M_6)_{\sigma_6 | \sigma_7} \dots$$

This is easy: both sides vanish if contracted with matrices d_i (just like in the previous examples, either due to nilpotency relations or due to complete antisymmetry of ϵ -tensors) and both sides are antisymmetric under permutations of indices inside sets σ_i . These constraints are very restrictive: any two solutions of these constraints must be proportional to each other.

The coefficient of proportionality is either equal to determinant of this complex or equal to inverse determinant of this complex, depending on the number of terms p . The easiest way to determine this sign is to notice, that the rightmost differential d_{p-1} should enter the expression for differential in numerator, not in denominator. Consequently,

$$\eta = \text{DET } (d_1, d_2, \dots, d_{p-1}) = \frac{(M_{p-1})_{\sigma_{p-1}} (M_{p-3})_{\sigma_{p-3} | \sigma_{p-2}} (M_{p-5})_{\sigma_{p-5} | \sigma_{p-4}} \dots}{(M_{p-2})_{\sigma_{p-2} | \sigma_{p-1}} (M_{p-4})_{\sigma_{p-4} | \sigma_{p-3}} (M_{p-6})_{\sigma_{p-6} | \sigma_{p-5}} \dots} \quad (3.3)$$

or, in a slightly more convenient form,

$$\eta^{(-1)^p} = \frac{(M_1)_{\sigma_2} (M_3)_{\sigma_3 | \sigma_4} (M_5)_{\sigma_5 | \sigma_6} \dots}{(M_2)_{\sigma_2 | \sigma_3} (M_4)_{\sigma_4 | \sigma_5} (M_6)_{\sigma_6 | \sigma_7} \dots}$$

By definition of σ_i , there are $C_{l_2}^{k_1} C_{l_3}^{k_2} \dots C_{l_{p-1}}^{k_{p-2}}$ different ways to select them. All these choices give one and the same determinant. This independence is ensured by the following facts:

1. Numerator and denominator depend on the same indices
2. Numerator and denominator are antisymmetric in each group of indices σ_i separately
3. Numerator and denominator satisfy one and the same highly overdefined system of linear constraints due to nilpotency of d and complete antisymmetry of ϵ

3.3 Degree of determinant

As follows from (3.3) and from the fact, that minor M_i has size k_i , resultant has degree

$$\deg \left(\text{DET } (d_1, d_2, \dots, d_{p-1}) \right) = \sum_{i=1}^{p-1} (-1)^{p+i+1} k_i = k_{p-1} - k_{p-2} + \dots$$

in coefficients of the differential operators. In terms of dimensions, we have

$$k_i = \sum_{j=1}^i (-1)^{i-j} l_j = l_i - l_{i-1} + \dots$$

$$\deg \left(\text{DET } (d_1, d_2, \dots, d_{p-1}) \right) = \sum_{i=1}^{p-1} i (-1)^{i-1} l_{p-i} = l_{p-1} - 2l_{p-2} + \dots \quad (3.4)$$

3.4 How to calculate determinant of complex: concise summary

For reference reasons and also for those, who need to calculate a determinant of their own complex and do not want to spend their time on reading the previous sections, in this section we give a short clear procedure how to calculate a determinant of arbitrary complex

$$L_1 \xrightarrow{d_1} L_2 \xrightarrow{d_2} \dots \xrightarrow{d_{p-1}} L_p$$

with vanishing Euler characteristic

$$\chi = \dim L_1 - \dim L_2 + \dim L_3 - \dots + (-1)^p \dim L_p = 0$$

Select a basis in each linear space, then complex is represented by a sequence of rectangular matrices. The $l_i \times l_{i+1}$ matrix d_i has l_i rows and l_{i+1} columns. The determinant of complex

$$\eta = \text{DET} (d_1, d_2, \dots, d_{p-1})$$

is constructed from minors of these matrices as follows. Take basis vectors in the linear space L_i as a set

$$\{1, 2, \dots, l_i\}$$

Select arbitrary subsets $\sigma_i \subset \{1, 2, \dots, l_i\}$ which consist of

$$|\sigma_i| = \sum_{k=1}^{i-1} (-1)^{i+k+1} l_k$$

elements, i.e.,

$$|\sigma_1| = 0$$

$$|\sigma_2| = l_1$$

$$|\sigma_3| = l_2 - l_1$$

$$|\sigma_4| = l_3 - l_2 + l_1$$

$$|\sigma_5| = l_4 - l_3 + l_2 - l_1$$

.....

Define conjugate subsets

$$\tilde{\sigma}_i = \{1, 2, \dots, l_i\} / \sigma_i$$

which consist of

$$|\tilde{\sigma}_i| = l_i - |\sigma_i| = |\sigma_{i+1}|$$

elements. Calculate minors

$$M_i = \text{the minor of } d_i, \text{ which occupies rows } \tilde{\sigma}_i \text{ and columns } \sigma_{i+1}$$

The determinant of complex equals

$$\boxed{\text{DET} (d_1, d_2, \dots, d_{p-1}) = \prod_{i=1}^{p-1} (M_i)^{(-1)^{p+i+1}} = \frac{M_{p-1} M_{p-3} \dots}{M_{p-2} M_{p-4} \dots}} \quad (3.5)$$

In fact, this answer does not depend on the choice of σ_i . More details are given in the previous section.

4 Koszul complexes

After discussing the general theory of complexes and their determinants, we return to Koszul complexes, relevant to the theory of resultants. The basic objects were already mentioned in the Introduction to the present paper.

Consider a polynomial map of type $n|r$

$$f : \begin{pmatrix} x_1 \\ x_2 \\ \dots \\ x_n \end{pmatrix} \mapsto \begin{pmatrix} f_1(x_1, x_2, \dots, x_n) \\ f_2(x_1, x_2, \dots, x_n) \\ \dots \\ f_n(x_1, x_2, \dots, x_n) \end{pmatrix}$$

i.e, of degree r in n variables:

$$f_i(x_1, x_2, \dots, x_n) = \sum_{j_1, \dots, j_r=1}^n f_i^{j_1, j_2, \dots, j_r} x_{j_1} x_{j_2} \dots x_{j_r}$$

Complement the commuting variables

$$x_1, x_2, \dots, x_n : x_i x_j - x_j x_i = 0$$

with anti-commuting variables

$$\theta_1, \theta_2, \dots, \theta_n : \theta_i \theta_j + \theta_j \theta_i = 0$$

and consider polynomials, depending both on $x_1, \dots, x_n$ and $\theta_1, \dots, \theta_n$. Denote through $\Omega(p, q)$ the space of such polynomials of degree p in x -variables and degree q in θ -variables. Their dimensions are

$$\dim \Omega(p, q) = C_{p+n-1}^p C_n^q \quad (4.1)$$

The degree q can not be greater than n , since θ are anti-commuting variables. *Koszul differential*, built from f , is a linear operator which acts on these spaces by the rule

$$\hat{d} = f_1(x_1, \dots, x_n) \frac{\partial}{\partial \theta_1} + \dots + f_n(x_1, \dots, x_n) \frac{\partial}{\partial \theta_n}$$

and is automatically nilpotent:

$$\hat{d}\hat{d} = f_j(x) f_k(x) \frac{\partial}{\partial \theta_j} \frac{\partial}{\partial \theta_k} = f_j(x) f_k(x) \left(\frac{\partial}{\partial \theta_j} \frac{\partial}{\partial \theta_k} + \frac{\partial}{\partial \theta_k} \frac{\partial}{\partial \theta_j} \right) = 0$$

It sends

$$\hat{d} : \Omega(p, q) \rightarrow \Omega(p+r, q-1)$$

giving rise to the following *Koszul complex*:

$$\Omega(p, q) \xrightarrow{\hat{d}} \Omega(p+r, q-1) \xrightarrow{\hat{d}} \Omega(p+2r, q-2) \xrightarrow{\hat{d}} \dots \xrightarrow{\hat{d}} \Omega(0, R)$$

Thus, for one and the same operator $\hat{d}$ there are many different Koszul complexes, depending on single integer parameter R – the x -degree of the rightmost space. Using the formula (4.1) for dimensions, it is easy to write down all of them. For example, the tower of Koszul complexes for the case $3|2$ is

R	Spaces	Dimensions	Euler characteristic
0	$\Omega(0, 0)$	1	1
1	$\Omega(1, 0)$	3	3
2	$\Omega(0, 1) \rightarrow \Omega(2, 0)$	$3 \rightarrow 6$	3
3	$\Omega(1, 1) \rightarrow \Omega(3, 0)$	$9 \rightarrow 10$	1
4	$\Omega(0, 2) \rightarrow \Omega(2, 1) \rightarrow \Omega(4, 0)$	$3 \rightarrow 18 \rightarrow 15$	0
5	$\Omega(1, 2) \rightarrow \Omega(3, 1) \rightarrow \Omega(5, 0)$	$9 \rightarrow 30 \rightarrow 21$	0
6	$\Omega(0, 3) \rightarrow \Omega(2, 2) \rightarrow \Omega(4, 1) \rightarrow \Omega(6, 0)$	$1 \rightarrow 18 \rightarrow 45 \rightarrow 28$	0
7	$\Omega(1, 3) \rightarrow \Omega(3, 2) \rightarrow \Omega(5, 1) \rightarrow \Omega(7, 0)$	$3 \rightarrow 30 \rightarrow 63 \rightarrow 36$	0
...	...	...	...

For reference reasons, let us write down several other examples explicitly:

	R	Dimensions	χ		R	Dimensions	χ
	0	1	1		0	1	1
	1	3	3		1	3	3
	2	6	6		2	6	6
	3	$3 \rightarrow 10$	7		3	10	10
	4	$9 \rightarrow 15$	6		4	$3 \rightarrow 15$	12
$\boxed{3 3}$	5	$18 \rightarrow 21$	3	$\boxed{3 4}$	5	$9 \rightarrow 21$	12
	6	$3 \rightarrow 30 \rightarrow 28$	1		6	$18 \rightarrow 28$	10
	7	$9 \rightarrow 45 \rightarrow 36$	0		7	$30 \rightarrow 36$	6
	8	$18 \rightarrow 63 \rightarrow 45$	0		8	$3 \rightarrow 45 \rightarrow 45$	3
	9	$1 \rightarrow 30 \rightarrow 84 \rightarrow 55$	0		9	$9 \rightarrow 63 \rightarrow 55$	1
	10	$3 \rightarrow 45 \rightarrow 108 \rightarrow 66$	0		10	$18 \rightarrow 84 \rightarrow 66$	0
	11	$6 \rightarrow 63 \rightarrow 135 \rightarrow 78$	0		11	$30 \rightarrow 108 \rightarrow 78$	0
	...	...	...		...	...	...

	R	Dimensions	χ
	0	1	1
	1	4	4
	2	$4 \rightarrow 10$	6
$\boxed{4 2}$	3	$16 \rightarrow 20$	4
	4	$6 \rightarrow 40 \rightarrow 35$	1
	5	$24 \rightarrow 80 \rightarrow 56$	0
	6	$4 \rightarrow 60 \rightarrow 140 \rightarrow 84$	0
	7	$16 \rightarrow 120 \rightarrow 224 \rightarrow 120$	0
	...	...	...

	R	Dimensions	χ		R	Dimensions	χ
	0	1	1		0	1	1
	1	4	4		1	4	4
	2	10	10		2	10	10
	3	$4 \rightarrow 20$	16		3	20	20
	4	$16 \rightarrow 35$	19		4	$4 \rightarrow 35$	31
	5	$40 \rightarrow 56$	16		5	$16 \rightarrow 56$	40
$\boxed{4 3}$	6	$6 \rightarrow 80 \rightarrow 84$	10	$\boxed{4 4}$	6	$40 \rightarrow 84$	44
	7	$24 \rightarrow 140 \rightarrow 120$	4		7	$80 \rightarrow 120$	40
	8	$60 \rightarrow 224 \rightarrow 165$	1		8	$6 \rightarrow 140 \rightarrow 165$	31
	9	$4 \rightarrow 120 \rightarrow 336 \rightarrow 220$	0		9	$24 \rightarrow 224 \rightarrow 220$	20
	10	$16 \rightarrow 210 \rightarrow 480 \rightarrow 286$	0		10	$60 \rightarrow 336 \rightarrow 286$	10
	11	$40 \rightarrow 336 \rightarrow 660 \rightarrow 364$	0		11	$120 \rightarrow 480 \rightarrow 364$	4
	12	$1 \rightarrow 80 \rightarrow 504 \rightarrow 880 \rightarrow 455$	0		12	$4 \rightarrow 210 \rightarrow 660 \rightarrow 455$	1
	13	$4 \rightarrow 140 \rightarrow 720 \rightarrow 1144 \rightarrow 560$	0		13	$16 \rightarrow 336 \rightarrow 880 \rightarrow 560$	0
	14	$10 \rightarrow 224 \rightarrow 990 \rightarrow 1456 \rightarrow 680$	0		14	$40 \rightarrow 504 \rightarrow 1144 \rightarrow 680$	0
	...	...	...		...	...	...

	R	Dimensions	χ
	0	1	1
	1	5	5
	2	$5 \rightarrow 15$	10
$\boxed{5 2}$	3	$25 \rightarrow 35$	10
	4	$10 \rightarrow 75 \rightarrow 70$	5
	5	$50 \rightarrow 175 \rightarrow 126$	1
	6	$10 \rightarrow 150 \rightarrow 350 \rightarrow 210$	0
	7	$50 \rightarrow 350 \rightarrow 630 \rightarrow 330$	0
	...	...	...

Many properties of Koszul complexes are evident already from these reference tables. For example, one can see that Euler characteristic χ vanishes, if $R > n(r-1)$. If it does, then, according to the previous section, the Koszul complex has a determinant

$$DET(d_1, \dots, d_{p-1})$$

where $p = 1 + [R/r]$ is the number of terms in the complex. The main fact, announced already in the title of this paper, is that determinant of Koszul complex is equal to the resultant of the original map:

$$R_{n|r}(f_1, \dots, f_n) = DET(d_1, \dots, d_{p-1}) \quad (4.2)$$

Note, that determinant of a complex is generally a rational function; however, Koszul complexes are very special complexes, and their determinants turn out to be polynomials (actually, the resultants of the underlying maps f). Despite the importance of this mathematical theorem, we do not include its proof into this text, it can be found, for example, in the book [5]. Instead of a proof, we give a number of illustrations and examples, which do not require any background in homological algebra.

The simplest way to understand the relation (4.2) is to notice, that determinant has the same degree in coefficients of $f_1, \dots, f_n$ as resultant. Indeed, substituting (4.1) into (3.4) we obtain

$$\begin{aligned} \deg \left(DET(d_1, d_2, \dots, d_{p-1}) \right) &= \sum_{i=1}^{p-1} i(-1)^{i-1} l_{p-i} = \sum_{i=1}^{[R/r]} i(-1)^{i-1} \dim \Omega(R-ir, i) = \\ &= \sum_{i=1}^{[R/r]} i(-1)^{i-1} C_n^i C_{R-ir+n-1}^{R-ir} = \sum_{i=1}^{[R/r]} \frac{(-1)^{i-1} n}{(n-i)!(i-1)!} \frac{(R-ir+n-1)!}{(R-ir)!} = \\ &= \sum_{i=1}^{[R/r]} \frac{(-1)^{i-1}}{(n-i)!(i-1)!} \oint \frac{dz}{z} z^{ir-R} \frac{n!}{(1-z)^n} = \\ &= \oint \frac{dz}{z} \frac{n!}{(1-z)^n} \cdot \sum_{i=1}^{[R/r]} \frac{(-1)^{i-1}}{(n-i)!(i-1)!} z^{ir-R} = \\ &= \oint \frac{dz}{z} \frac{n!}{(1-z)^n} \cdot \frac{z^{r-R}(1-z^r)^{n-1}}{(n-1)!} = \\ &= n \oint \frac{dz(1-z^r)^{n-1}}{(1-z)^n} z^{r-R-1} = n \lim_{z \rightarrow 1} \left(\frac{1-z^r}{1-z} \right)^{n-1} = nr^{n-1} \end{aligned}$$

i.e. if determinant is defined, its degree equals to degree (2.3) of the resultant:

$$\deg \left(DET(d_1, d_2, \dots, d_{p-1}) \right) = \deg \left(R_{n|r}(f_1, \dots, f_n) \right)$$

Consequently,

$$DET(d_1, d_2, \dots, d_{p-1}) = C(d_1, d_2, \dots, d_{p-1}) \cdot R_{n|r}(f_1, \dots, f_n)$$

where the coefficient of proportionality C has degree 0. Since determinant of Koszul complex is polynomial rather than a rational function [5], C is just a constant and does not depend on anything. This justifies (4.2).

The second, even more direct, way to establish the relation between resultant and determinant of Koszul complex, is to consider the rightmost differential in the Koszul complex:

$$\hat{d}: \Omega(R-r, 1) \xrightarrow{\hat{d}} \Omega(R, 0)$$

which acts on these spaces as

$$(x_1^{i_1} x_2^{i_2} \dots x_n^{i_n}) \cdot \theta_j \mapsto (x_1^{i_1} x_2^{i_2} \dots x_n^{i_n}) \cdot f_j(x_1, \dots, x_n)$$

and is represented by $nC_{R-r+n-1}^{R-r} \times C_{R+n-1}^R$ matrix d_{p-1} . If resultant $R_{n|r}(f_1, \dots, f_n)$ vanishes, then

$$\begin{cases} f_1(x_1, x_2, \dots, x_n) = 0 \\ f_2(x_1, x_2, \dots, x_n) = 0 \\ \dots \\ f_n(x_1, x_2, \dots, x_n) = 0 \end{cases}$$

becomes solvable, i.e. a vector $(v_1, \dots, v_n)$ exists, such that all $f_j(v_1, \dots, v_n) = 0$. As a consequence, there is a $nC_{R-r+n-1}^{R-r}$ -dimensional vector

$$(v_1^{i_1} v_2^{i_2} \dots v_n^{i_n}) \cdot \theta_j$$

from $\Omega(R-r, 1)$, annihilated by d_{p-1} . Therefore,

$$R_{n|r} = 0 \Rightarrow \text{rank of } d_{p-1} \text{ drops down by one}$$

what is the same,

$$R_{n|r} = 0 \Rightarrow \text{all top-dimensional minors of } d_{p-1} \text{ vanish}$$

This is only possible if all top-dimensional minors of d_{p-1} are divisible by $R_{n|r}$. As follows from (3.3), determinant of such complex is also divisible by $R_{n|r}$. Whenever resultant $R_{n|r}$ vanishes, determinant of Koszul complex also vanishes. Again, this is enough to state (4.2), if determinant of Koszul complex is known to be a polynomial.

To finish our discussion of Koszul complexes, let us derive a simple generating function

$$\sum_{R=0}^{\infty} \chi_R t^R = \left(\frac{1-t^r}{1-t} \right)^n \quad (4.3)$$

for their Euler characteristics χ_R . They are quite interesting: for $r = 2$, they coincide with binomial coefficients (as one can see in the reference tables), thus for $r > 2$ they provide a generalization of binomial coefficients. Using exactly the same contour integral trick, we obtain

$$\begin{aligned} \sum_{R=0}^{\infty} \chi_R t^R &= \sum_{R=0}^{\infty} \sum_{i=0}^{\infty} (-1)^i \dim \Omega(R-ir, i) t^R = \sum_{R=0}^{\infty} \sum_{i=0}^{\infty} (-1)^i C_n^i C_{R-ir+n-1}^{R-ir} t^R = \\ &= \sum_{R=0}^{\infty} \sum_{i=0}^{\infty} \frac{(-1)^i n!}{i!(n-i)!} \frac{(R-ir+n-1)!}{(R-ir)!} t^R = \sum_{R=0}^{\infty} \sum_{i=0}^{\infty} \frac{(-1)^i}{i!(n-i)!} t^R \oint \frac{dz}{z} z^{ir-R} \frac{n!}{(1-z)^n} = \\ &= \oint \frac{dz}{z} \frac{n!}{(1-z)^n} \cdot \sum_{R=0}^{\infty} \sum_{i=0}^{\infty} \frac{(-1)^i}{i!(n-i)!} t^R z^{ir-R} = \\ &= \oint \frac{dz}{z} \frac{n!}{(1-z)^n} \cdot \frac{z(1-z^r)^n}{n!(z-t)} = \\ &= \oint \frac{dz}{z-t} \left(\frac{1-z^r}{1-z} \right)^n = \left(\frac{1-t^r}{1-t} \right)^n \end{aligned}$$

and the generating function (4.3) is derived.

5 Evaluation of $R_{3|2}$

As a final illustration to the whole paper, let us consider in detail the case $3|2$, a system of three quadratic equations in three unknowns:

$$\begin{cases} F_1 = a_{11}x_1^2 + a_{12}x_1x_2 + a_{13}x_1x_3 + a_{22}x_2^2 + a_{23}x_2x_3 + a_{33}x_3^2 \\ F_2 = b_{11}x_1^2 + b_{12}x_1x_2 + b_{13}x_1x_3 + b_{22}x_2^2 + b_{23}x_2x_3 + b_{33}x_3^2 \\ F_3 = c_{11}x_1^2 + c_{12}x_1x_2 + c_{13}x_1x_3 + c_{22}x_2^2 + c_{23}x_2x_3 + c_{33}x_3^2 \end{cases}$$

This case is very important in resultant theory, for a number of reasons. First of all, it is the simplest example of multidimensional ($n > 2$) resultants. At the same time, it is quite a representative example. $R_{3|2}(F_1, F_2, F_3)$ is a homogeneous polynomial of degree $3 \cdot 2^2 = 12$ in a_{ij}, b_{ij}, c_{ij} . If completely expanded, this polynomial consists of 21894 monomials and takes 30-50 pages of A4 paper, depending on the font, to write it down. It is quite amazing, that such a simple system of equations has such an enormously lengthy consistency (solvability) condition. Today we know, that these 21894 terms are in fact controlled by various hidden structures [6, 21]. Explicit knowledge of these structures allows to write down $R_{3|2}$ in a reasonably short form. Koszul complex, which represents $R_{3|2}$ through several matrices of small size, is historically the first example of such a structure.

As usual, Koszul differential is

$$\hat{d} = F_1 \frac{\partial}{\partial \theta_1} + F_2 \frac{\partial}{\partial \theta_2} + F_3 \frac{\partial}{\partial \theta_3}$$

The tower of Koszul complexes is

R	Spaces	Dimensions	Euler characteristic
0	$\Omega(0, 0)$	1	1
1	$\Omega(1, 0)$	3	3
2	$\Omega(0, 1) \rightarrow \Omega(2, 0)$	$3 \rightarrow 6$	3
3	$\Omega(1, 1) \rightarrow \Omega(3, 0)$	$9 \rightarrow 10$	1
4	$\Omega(0, 2) \rightarrow \Omega(2, 1) \rightarrow \Omega(4, 0)$	$3 \rightarrow 18 \rightarrow 15$	0
5	$\Omega(1, 2) \rightarrow \Omega(3, 1) \rightarrow \Omega(5, 0)$	$9 \rightarrow 30 \rightarrow 21$	0
6	$\Omega(0, 3) \rightarrow \Omega(2, 2) \rightarrow \Omega(4, 1) \rightarrow \Omega(6, 0)$	$1 \rightarrow 18 \rightarrow 45 \rightarrow 28$	0
7	$\Omega(1, 3) \rightarrow \Omega(3, 2) \rightarrow \Omega(5, 1) \rightarrow \Omega(7, 0)$	$3 \rightarrow 30 \rightarrow 63 \rightarrow 36$	0
...	...	...	...

Let us describe in detail the simplest case $R = 4$ and next-to-the-simplest case $R = 5$. The complex with $R = 4$, is the first determinant-possessing complex (i.e. has a vanishing Euler characteristics):

$$\Omega(0, 2) \rightarrow \Omega(2, 1) \rightarrow \Omega(4, 0)$$

with dimensions $3 \rightarrow 18 \rightarrow 15$. If we select a basis in $\Omega(0, 2)$ as

$$\{\theta_2\theta_3, \theta_1\theta_3, \theta_1\theta_2\}$$

a basis in $\Omega(2, 1)$ as

$$\begin{aligned} &\{x_1^2\theta_3, x_1^2\theta_2, x_1^2\theta_1, x_1x_2\theta_3, x_1x_2\theta_2, x_1x_2\theta_1, x_1x_3\theta_3, x_1x_3\theta_2, \\ &x_1x_3\theta_1, x_2^2\theta_3, x_2^2\theta_2, x_2^2\theta_1, x_2x_3\theta_3, x_2x_3\theta_2, x_2x_3\theta_1, x_3^2\theta_3, x_3^2\theta_2, x_3^2\theta_1\} \end{aligned}$$

and, finally, a basis in $\Omega(4, 0)$ as

$$\{x_1^4, x_1^3x_2, x_1^3x_3, x_1^2x_2^2, x_1^2x_2x_3, x_1^2x_3^2, x_1x_2^3, x_1x_2^2x_3, x_1x_2x_3^2, x_1x_3^3, x_2^4, x_2^3x_3, x_2^2x_3^2, x_2x_3^3, x_3^4\}$$

then Koszul differential is represented by a pair of matrices: one 3×18

$$d_1 = \begin{bmatrix} b_{1,1} & c_{1,1} & 0 & b_{1,2} & c_{1,2} & 0 & b_{1,3} & c_{1,3} & 0 & b_{2,2} & c_{2,2} & 0 & b_{2,3} & c_{2,3} & 0 & b_{3,3} & c_{3,3} & 0 \\ -a_{1,1} & 0 & c_{1,1} & -a_{1,2} & 0 & c_{1,2} & -a_{1,3} & 0 & c_{1,3} & -a_{2,2} & 0 & c_{2,2} & -a_{2,3} & 0 & c_{2,3} & -a_{3,3} & 0 & c_{3,3} \\ 0 & -a_{1,1} & -b_{1,1} & 0 & -a_{1,2} & -b_{1,2} & 0 & -a_{1,3} & -b_{1,3} & 0 & -a_{2,2} & -b_{2,2} & 0 & -a_{2,3} & -b_{2,3} & 0 & -a_{3,3} & -b_{3,3} \end{bmatrix}$$

and another 18×15

$$d_2 = \begin{bmatrix} c_{1,1} & c_{1,2} & c_{1,3} & c_{2,2} & c_{2,3} & c_{3,3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -b_{1,1} & -b_{1,2} & -b_{1,3} & -b_{2,2} & -b_{2,3} & -b_{3,3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ a_{1,1} & a_{1,2} & a_{1,3} & a_{2,2} & a_{2,3} & a_{3,3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & c_{1,1} & 0 & c_{1,2} & c_{1,3} & 0 & c_{2,2} & c_{2,3} & c_{3,3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -b_{1,1} & 0 & -b_{1,2} & -b_{1,3} & 0 & -b_{2,2} & -b_{2,3} & -b_{3,3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & a_{1,1} & 0 & a_{1,2} & a_{1,3} & 0 & a_{2,2} & a_{2,3} & a_{3,3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & c_{1,1} & 0 & c_{1,2} & c_{1,3} & 0 & c_{2,2} & c_{2,3} & c_{3,3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -b_{1,1} & 0 & -b_{1,2} & -b_{1,3} & 0 & -b_{2,2} & -b_{2,3} & -b_{3,3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & a_{1,1} & 0 & a_{1,2} & a_{1,3} & 0 & a_{2,2} & a_{2,3} & a_{3,3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & c_{1,1} & 0 & 0 & c_{1,2} & c_{1,3} & 0 & 0 & c_{2,2} & c_{2,3} & c_{3,3} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -b_{1,1} & 0 & 0 & -b_{1,2} & -b_{1,3} & 0 & 0 & -b_{2,2} & -b_{2,3} & -b_{3,3} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & a_{1,1} & 0 & 0 & a_{1,2} & a_{1,3} & 0 & 0 & a_{2,2} & a_{2,3} & a_{3,3} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & c_{1,1} & 0 & 0 & c_{1,2} & c_{1,3} & 0 & 0 & c_{2,2} & c_{2,3} & c_{3,3} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -b_{1,1} & 0 & 0 & -b_{1,2} & -b_{1,3} & 0 & 0 & -b_{2,2} & -b_{2,3} & -b_{3,3} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & a_{1,1} & 0 & 0 & a_{1,2} & a_{1,3} & 0 & 0 & a_{2,2} & a_{2,3} & a_{3,3} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & c_{1,1} & 0 & 0 & c_{1,2} & c_{1,3} & 0 & 0 & c_{2,2} & c_{2,3} & c_{3,3} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -b_{1,1} & 0 & 0 & -b_{1,2} & -b_{1,3} & 0 & 0 & -b_{2,2} & -b_{2,3} & -b_{3,3} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & a_{1,1} & 0 & 0 & a_{1,2} & a_{1,3} & 0 & 0 & a_{2,2} & a_{2,3} & a_{3,3} & 0 & 0 \end{bmatrix}$$

By selecting some 3 columns in $\hat{d}_1$ and complementary 15 rows in $\hat{d}_2$, we obtain the desired resultant

$$R = \begin{bmatrix} a_{1,1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ a_{1,2} & c_{1,1} & a_{1,1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ a_{1,3} & 0 & 0 & c_{1,1} & -b_{1,1} & a_{1,1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ a_{2,2} & c_{1,2} & a_{1,2} & 0 & 0 & 0 & c_{1,1} & -b_{1,1} & a_{1,1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ a_{2,3} & c_{1,3} & a_{1,3} & c_{1,2} & -b_{1,2} & a_{1,2} & 0 & 0 & 0 & c_{1,1} & -b_{1,1} & a_{1,1} & 0 & 0 & 0 & 0 \\ a_{3,3} & 0 & 0 & c_{1,3} & -b_{1,3} & a_{1,3} & 0 & 0 & 0 & 0 & 0 & 0 & c_{1,1} & -b_{1,1} & a_{1,1} & 0 \\ 0 & c_{2,2} & a_{2,2} & 0 & 0 & 0 & c_{1,2} & -b_{1,2} & a_{1,2} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & c_{2,3} & a_{2,3} & c_{2,2} & -b_{2,2} & a_{2,2} & c_{1,3} & -b_{1,3} & a_{1,3} & c_{1,2} & -b_{1,2} & a_{1,2} & 0 & 0 & 0 & 0 \\ 0 & c_{3,3} & a_{3,3} & c_{2,3} & -b_{2,3} & a_{2,3} & 0 & 0 & 0 & c_{1,3} & -b_{1,3} & a_{1,3} & c_{1,2} & -b_{1,2} & a_{1,2} & 0 \\ 0 & 0 & 0 & c_{3,3} & -b_{3,3} & a_{3,3} & 0 & 0 & 0 & 0 & 0 & 0 & c_{1,3} & -b_{1,3} & a_{1,3} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & c_{2,2} & -b_{2,2} & a_{2,2} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & c_{2,3} & -b_{2,3} & a_{2,3} & c_{2,2} & -b_{2,2} & a_{2,2} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & c_{3,3} & -b_{3,3} & a_{3,3} & c_{2,3} & -b_{2,3} & a_{2,3} & c_{2,2} & -b_{2,2} & a_{2,2} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & c_{3,3} & -b_{3,3} & a_{3,3} & c_{2,3} & -b_{2,3} & a_{2,3} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & c_{3,3} & -b_{3,3} & a_{3,3} & 0 \end{bmatrix}$$

$$\begin{bmatrix} b_{1,1} & c_{1,1} & c_{1,2} \\ -a_{1,1} & 0 & 0 \\ 0 & -a_{1,1} & -a_{1,2} \end{bmatrix}$$

This particular formula corresponds to the choice of columns 1, 2 and 5 in the first matrix, but the resultant, of course, does not depend on this choice: any other three columns will do. One only needs to care that determinant in the denominator does not vanish. If it is non-zero, then the upper minor is divisible by the lower minor, and the fraction is equal (up to sign) to the resultant.

Alternative complex is the second complex with vanishing χ , that corresponds to $R = 5$:

$$\Omega(1, 2) \rightarrow \Omega(3, 1) \rightarrow \Omega(5, 0)$$

It has dimensions $9 \rightarrow 30 \rightarrow 21$. If we select a basis in $\Omega(1, 2)$ as

$$\{x_1\theta_2\theta_3, x_1\theta_1\theta_3, x_1\theta_1\theta_2, x_2\theta_2\theta_3, x_2\theta_1\theta_3, x_2\theta_1\theta_2, x_3\theta_2\theta_3, x_3\theta_1\theta_3, x_3\theta_1\theta_2\}$$

a basis in $\Omega(3, 1)$ as

$$\begin{aligned} & \{x_1^3\theta_3, x_1^3\theta_2, x_1^3\theta_1, x_1^2x_2\theta_3, x_1^2x_2\theta_2, x_1^2x_2\theta_1, x_1^2x_3\theta_3, x_1^2x_3\theta_2, x_1^2x_3\theta_1, \\ & x_1x_2^2\theta_3, x_1x_2^2\theta_2, x_1x_2^2\theta_1, x_1x_2x_3\theta_3, x_1x_2x_3\theta_2, x_1x_2x_3\theta_1, x_1x_3^2\theta_3, x_1x_3^2\theta_2, x_1x_3^2\theta_1, x_2^3\theta_3, x_2^3\theta_2, x_2^3\theta_1, \\ & x_2^2x_3\theta_3, x_2^2x_3\theta_2, x_2^2x_3\theta_1, x_2x_3^2\theta_3, x_2x_3^2\theta_2, x_2x_3^2\theta_1, x_3^3\theta_3, x_3^3\theta_2, x_3^3\theta_1 \\ & x_1x_3\theta_1, x_2^2\theta_3, x_2^2\theta_2, x_2^2\theta_1, x_2x_3\theta_3, x_2x_3\theta_2, x_2x_3\theta_1, x_3^2\theta_3, x_3^2\theta_2, x_3^2\theta_1\} \end{aligned}$$

and, finally, a basis in $\Omega(5, 0)$ as

$$\{x_1^5, x_1^4x_2, x_1^4x_3, x_1^3x_2^2, x_1^3x_2x_3, x_1^3x_3^2, x_1^2x_2^3, x_1^2x_2^2x_3, x_1^2x_2x_3^2, x_1^2x_3^3, \\ x_1x_2^4, x_1x_2^3x_3, x_1x_2^2x_3^2, x_1x_2x_3^3, x_1x_3^4, x_2^5, x_2^4x_3, x_2^3x_3^2, x_2^2x_3^3, x_2x_3^4, x_3^5\}$$

then Koszul differential is represented by a pair of matrices: one 9×30

[illegible]

and another 30×21

[illegible]

By selecting some 9 columns in $\hat{d}_1$ and complementary 21 rows in $\hat{d}_2$, we obtain the desired resultant

$$R = \begin{bmatrix} -\partial_{1,1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -\partial_{1,2} & \epsilon_{1,1} & a_{1,1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -\partial_{1,3} & 0 & 0 & -\partial_{1,1} & a_{1,1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -\partial_{2,2} & \epsilon_{1,2} & a_{1,2} & 0 & 0 & \epsilon_{1,1} & -\partial_{1,1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -\partial_{2,3} & \epsilon_{1,3} & a_{1,3} & -\partial_{1,2} & a_{1,2} & 0 & 0 & \epsilon_{1,1} & a_{1,1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -\partial_{3,3} & 0 & 0 & -\partial_{1,3} & a_{1,3} & 0 & 0 & 0 & 0 & -\partial_{1,1} & a_{1,1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \epsilon_{2,2} & a_{2,2} & 0 & 0 & \epsilon_{1,2} & -\partial_{1,2} & 0 & 0 & 0 & 0 & -\partial_{1,1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \epsilon_{2,3} & a_{2,3} & -\partial_{2,2} & a_{2,2} & \epsilon_{1,3} & -\partial_{1,3} & \epsilon_{1,2} & a_{1,2} & 0 & 0 & 0 & \epsilon_{1,1} & -\partial_{1,1} & a_{1,1} & 0 & 0 & 0 & 0 & 0 \\ 0 & \epsilon_{3,3} & a_{3,3} & -\partial_{2,3} & a_{2,3} & 0 & 0 & \epsilon_{1,3} & a_{1,3} & -\partial_{1,2} & a_{1,2} & 0 & 0 & 0 & 0 & \epsilon_{1,1} & -\partial_{1,1} & a_{1,1} & 0 & 0 \\ 0 & 0 & 0 & -\partial_{3,3} & a_{3,3} & 0 & 0 & 0 & 0 & -\partial_{1,3} & a_{1,3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \epsilon_{1,1} & -\partial_{1,1} & a_{1,1} \\ 0 & 0 & 0 & 0 & 0 & \epsilon_{2,2} & -\partial_{2,2} & 0 & 0 & 0 & 0 & -\partial_{1,2} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \epsilon_{2,3} & -\partial_{2,3} & \epsilon_{2,2} & a_{2,2} & 0 & 0 & -\partial_{1,3} & \epsilon_{1,2} & -\partial_{1,2} & a_{1,2} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \epsilon_{3,3} & -\partial_{3,3} & \epsilon_{2,3} & a_{2,3} & -\partial_{2,2} & a_{2,2} & 0 & \epsilon_{1,3} & -\partial_{1,3} & a_{1,3} & \epsilon_{1,2} & -\partial_{1,2} & a_{1,2} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \epsilon_{3,3} & a_{3,3} & -\partial_{2,3} & a_{2,3} & 0 & 0 & 0 & 0 & \epsilon_{1,3} & -\partial_{1,3} & a_{1,3} & \epsilon_{1,2} & -\partial_{1,2} & a_{1,2} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -\partial_{3,3} & a_{3,3} & 0 & 0 & 0 & 0 & 0 & 0 & \epsilon_{1,3} & -\partial_{1,3} & a_{1,3} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -\partial_{2,2} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -\partial_{2,3} & \epsilon_{2,2} & -\partial_{2,2} & a_{2,2} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -\partial_{3,3} & \epsilon_{2,3} & -\partial_{2,3} & a_{2,3} & \epsilon_{2,2} & -\partial_{2,2} & a_{2,2} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \epsilon_{3,3} & -\partial_{3,3} & a_{3,3} & \epsilon_{2,3} & -\partial_{2,3} & a_{2,3} & \epsilon_{2,2} & -\partial_{2,2} & a_{2,2} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \epsilon_{3,3} & -\partial_{3,3} & a_{3,3} & \epsilon_{2,3} & -\partial_{2,3} & a_{2,3} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \epsilon_{3,3} & -\partial_{3,3} & a_{3,3} \end{bmatrix}$$

$$= \begin{bmatrix} \partial_{1,1} & 0 & \epsilon_{1,2} & \partial_{1,3} & 0 & \epsilon_{2,3} & \partial_{3,3} & 0 & 0 \\ -a_{1,1} & \epsilon_{1,1} & 0 & -a_{1,3} & \epsilon_{2,2} & 0 & -a_{3,3} & 0 & 0 \\ 0 & -\partial_{1,1} & -a_{1,2} & 0 & -\partial_{2,2} & -a_{2,3} & 0 & 0 & 0 \\ 0 & 0 & \epsilon_{1,1} & 0 & 0 & \epsilon_{1,3} & 0 & \partial_{2,2} & 0 \\ 0 & 0 & 0 & 0 & \epsilon_{1,2} & 0 & 0 & -a_{2,2} & \epsilon_{2,2} \\ 0 & 0 & -a_{1,1} & 0 & -\partial_{1,2} & -a_{1,3} & 0 & 0 & -\partial_{2,2} \\ 0 & 0 & 0 & \partial_{1,1} & 0 & \epsilon_{1,2} & \partial_{1,3} & 0 & 0 \\ 0 & 0 & 0 & -a_{1,1} & 0 & 0 & -a_{1,3} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -a_{1,2} & 0 & 0 & 0 \end{bmatrix}$$

This particular formula corresponds to the choice of columns 1, 3, 5, 7, 12, 14, 16, 19 and 21 in the first matrix. Of course, any other choice will give the the same (up to a sign) result, if only the minor in denominator is non-zero. Direct calculation (division of one minor by another) should demonstrate, that the two complexes $3 \rightarrow 18 \rightarrow 15$ and $9 \rightarrow 30 \rightarrow 21$ give one and the same expression for the resultant. It should also be reproduced by determinants of all other complexes from the $3|2$ list with $R > 3$.

There are alternative representations, providing $R_{3|2}$ in far more compact form, see, for example, [6, 16, 21]. Still, advantage of Koszul complexes is that they provide an explicit answer for *any* resultant by a straightforward, generic and simply-formulated algorithm:

$$\text{Resultant of a map} = \text{Determinant of any associated Koszul complex} \quad (5.1)$$

This answer is excessively complicated in the following two senses: (i) the ratio of various determinants at the r.h.s. is actually reduced to a polynomial, but this cancelation is not explicit in (5.1); (ii) the r.h.s. is actually independent of the choice of Koszul complex, but this independence is not explicit in (5.1). However, until powerful alternatives were recently found in [11, 12, 16], relation (5.1) remained for some years the only *universal* representation for generic resultants.

6 Conclusion

This paper is a brief review of three important notions of tomorrow's mathematical physics: determinant of linear complex, resultant of non-linear map and Koszul complex. Each of them has its own value and large domain of potential applications. Remarkably, they are interrelated: resultant is a determinant of a Koszul complex. We gave a number of simple illustrations to this general theorem which do not require any preliminary knowledge, but reveal all essential nuances and ambiguities. Still further simplification of this presentation would be important for making these technical means into a nicely developed and widely used tool of theoretical physics and string theory.

Acknowledgements

We are indebted for illuminating explanations and discussions to V.Dolotin and A.Gorodentsev. Our work is partly supported by Russian Federal Nuclear Energy Agency and the Russian President's Grant of Support for the Scientific Schools NSh-3036.2008.2, by RFBR grants 07-01-00526 (A.A.) and 07-02-00645 (A.M. & Sh.Sh.), by the joint grants 09-01-92440-CE and 09-02-91005-ANF and by the NWO project 047.011.2004.026 (A.M.). The work of Sh.Shakirov is also supported in part by the Moebius Contest Foundation for Young Scientists and by the Dynasty Foundation.

References

- [1] A. Jennings, *Matrices, ancient and modern*, Bull. Inst. Math. Appl. **13** (5) (1977) 117 – 123
- [2] E. Forbes, *The astronomical work of Carl Friedrich Gauss (1777-1855)*, Historia Math. **5** (2) (1978) 167 – 181
- [3] A. Cayley, *On the theory of elimination*, Cambridge and Dublin Mathematical J. **3** (1848) 116 - 120
F.S. Macaulay, *On some Formulae in Elimination*, Proceedings of The London Mathematical Society, Vol. **XXXV** (1903) 3 - 27
A.L. Dixon, *The eliminant of three quantics in two independent variables*, Proceedings of The London Mathematical Society, **6** (1908) 468 - 478
E. Bézout, *Théorie générale des Equations Algébriques*, 1779, Paris
- [4] J.J. Sylvester, *On a general method of determining by mere inspection the derivations from two equations of any degree*, Philosophical Magazine **16** (1840) 132 - 135
- [5] I. Gelfand, M. Kapranov, and A. Zelevinsky, *Discriminants, Resultants and Multidimensional Determinants*, Birkhauser, 1994
- [6] V. Dolotin and A. Morozov, *Introduction to Non-Linear Algebra*, World Scientific, 2007; arXiv:hep-th/0609022
- [7] A. Miyake and M. Wadati, *Multiparticle Entanglement and Hyperdeterminants*, arXiv:quant-ph/02121146;
M. Duff, *String Triality, Black Hole Entropy and Cayleyes Hyperdeterminant*, arXiv:hep-th/0601134;
Hidden Symmetries of the Nambu-Goto Action, arXiv:hep-th/0602160;
A. Linde and R. Kallosh, *Strings, Black Holes and Quantum Information*, arXiv:hep-th/0602061
- [8] S. Kachru, A. Klemm, W. Lerche, P. Mayr and C. Vafa, *Nonperturbative Results on the Point Particle Limit of N=2 Heterotic String Compactifications*, Nucl.Phys. **B459** (1996) 537-558, arXiv:hep-th/9508155;
T. Eguchi and Y. Tachikawa, *Rigid Limit in N=2 Supergravity and Weak-Gravity Conjecture*, arXiv:hep-th/0706.2114;
P. Aspinwall, B. Greene and d. Morrison, *Measuring Small Distances in N=2 Sigma Models*, Nucl.Phys. **B420** (1994) 184-242, arXiv:hep-th/9311042

- [9] A. Morozov and A. Niemi, *Can Renormalization Group Flow End in a Big Mess?* Nucl.Phys. **B666** (2003) 311-336, arXiv:hep-th/0304178;
V. Dolotin and A. Morozov, *The Universal Mandelbrot Set. Beginning of the Story*, arXiv:hep-th/0501235; Int.J.Mod.Phys. **A23** (2008) 3613-3684, arXiv:hep-th/0701234;
An. Morozov, *Universal Mandelbrot Set as a Model of Phase Transition Theory*, JETP Lett. **86** (2007) 745-748, arXiv:nlin/0710.2315
- [10] D. Manocha, *Algebraic and Numeric Techniques for Modeling and Robotics*, PhD thesis, Computer Science Division, Department of Electrical Engineering and Computer Science, University of California, Berkeley.
- [11] A. Morozov and Sh. Shakirov, *Resultants and Contour Integrals*, arXiv:0807.4539
- [12] B. Gustafsson and V. Tkachev, *The resultant on compact Riemann surfaces*, arXiv:math/0710.2326, to appear in Comm.Math.Phys
- [13] E. Witten, *Topological Sigma Models*, Commun. Math. Phys. **118** (1988) 411;
E. Witten, *Mirror Manifolds And Topological Field Theory*, hep-th/9112056;
A.Losev, I.Polyubin, *On Connection between Topological Landau-Ginzburg Gravity and Integrable Systems*, hep-th/9305079;
M. Alexandrov, M. Kontsevich, A. Schwarz and O. Zaboronsky, *The Geometry of the master equation and topological quantum field theory*, Int. J. Mod. Phys. A **12** (1997) 1405; hep-th/9502010;
A. Losev, N. Nekrasov, S. Shatashvili, *Issues in Topological Gauge Theory*, hep-th/9711108
- [14] L.D. Faddeev and V.N. Popov, Phys. Lett. **B 25** (1967) 29;
I. V. Tyutin, *Gauge Invariance In Field Theory And Statistical Physics In Operator Formalism*, LEBEDEV-75-39;
R.Stora, *Progress in Gauge Field Theory* New York: Plenum Press, 1984;
I. Batalin and G. Vilkoviski, Nucl. Phys. **B234** (1984) 106;
A. Schwarz, *Geometry of Batalin-Vilkovisky quantization*, Commun. Math. Phys. **155** (1993) 249; hep-th/9205088
- [15] M. Douglas, *D-branes, Categories and N=1 Supersymmetry*, J.Math.Phys. **42** (2001) 2818-2843, arXiv:hep-th/0011017;
G. Moore, *Some Comments on Branes, G-flux, and K-theory*, Int.J.Mod.Phys. **A16** (2001) 936-944;
C.I. Lazaroiu, *Unitarity, D-brane dynamics and D-brane categories*, JHEP **0112** (2001) 031, arXiv:hep-th/0102183;
D.E. Diaconescu, *Enhanced D-Brane Categories from String Field Theory*, JHEP **0106** (2001) 016, arXiv:hep-th/0104200;
A. Kapustin and D. Orlov, *Remarks on A-branes, Mirror Symmetry, and the Fukaya category*, J.Geom.Phys. **48** (2003) 84, arXiv:hep-th/0109098;
J. Distler, H. Jockers and H. Park, *D-Brane Monodromies, Derived Categories and Boundary Linear Sigma Models*, arXiv:hep-th/0206242;
Sh. Katz and E. Sharpe, *D-branes, open string vertex operators, and Ext groups*, Adv.Theor.Math.Phys. **6** (2003) 979-1030, arXiv:hep-th/0208104;
P.S. Aspinwall, *D-Branes on Calabi-Yau Manifolds*, hep-th/0403166;
O. Lechtenfeld, A.D. Popov and R.J. Szabo, *Rank Two Quiver Gauge Theory, Graded Connections and Noncommutative Vortices*, JHEP **0609** (2006) 054, arXiv:hep-th/0603232;
A. Kapustin and E. Witten, *Electric-Magnetic Duality And The Geometric Langlands Program*, arXiv:hep-th/0604151
- [16] A. Morozov and Sh. Shakirov, *Analogue of the identity $\text{Log Det} = \text{Trace Log}$ for resultants*, arXiv:math-ph/0804.4632
- [17] D. Eisenbud and F.O. Schreyer, *Resultants and Chow forms via exterior syzygies*, J. Amer. Math. Soc. **16** (2003) 537-579, arXiv:math/0111040
- [18] C. Andrea, A.Dickenstein, *Explicit formulas for the multivariate resultant*, arXiv:math/0007036;
M. Chardin. *Formules à la Macaulay pour les sous-resultants en plusieurs variables*, C. R. Acad. Sci.

- Paris, **319** (1994) 433 – 436;
d. Manocha and J.Canny, *Multipolynomial Resultant Algorithms*, J.Symb.Comp. **15** (1993) 99 – 122;
J. Canny, E.Kaltofen and L.Yagati, *Solving Systems of Non-Linear Equations Faster*,
Proc.Internat.Symp.Symbolic Algebraic Comput., (1989) 121-128
- [19] A. Ostrowski. *The irreducibility of the resultant and connected irreducibility theorems*. **29** (1977) 252-260
L. Buse and C. D’Andrea, *On the irreducibility of multivariate subresultants*, Compt.Rend.Math. **338(4)** (2004) 287-290, arXiv:math/0309374
- [20] A. Kostrikin, *Introduction to algebra*, Springer-Verlag, New York, 1982
- [21] A. Morozov and Sh. Shakirov, *Introduction to Integral Discriminants*, to appear